

Package ‘SeedImbibition’

July 21, 2025

Type Package

Title Seed Imbibition Percentage

Version 0.1.0

Maintainer Tanuj Misra <tanujmisra102@gmail.com>

Description

Imbibition causes seeds to expand, which results in the seed coat or testa being broken. Seed germination begins with imbibition. Imbibition aids in the transport of water into the developing ovules. Imbibition is required during the first stages of root water absorption.

License GPL-3

Encoding UTF-8

NeedsCompilation no

Author Sudip Kumar Dutta [aut, cph],
Tanuj Misra [aut, cre, cph]

Repository CRAN

Date/Publication 2024-05-31 14:32:52 UTC

Contents

SeedImbibition	1
Index	3

SeedImbibition	<i>Seed Imbibition Percentage</i>
----------------	-----------------------------------

Description

Imbibition causes seeds to expand, which results in the seed coat or testa being broken. Seed germination begins with imbibition. Imbibition aids in the transport of water into the developing ovules. Imbibition is required during the first stages of root water absorption. Imbibition percentage of seeds was calculated using the following formula (Govender et al., 2008) <doi:10.1016/j.sajb.2007.10.006>

Usage

```
seed_imbib(wt_of_seed_after_imbib, initial_wt_of_seed)
```

Arguments

```
wt_of_seed_after_imbib  
    Weight of seeds after imbibition  
initial_wt_of_seed  
    Initial weight of seeds
```

Value

Seed Imbibition Percentage

References

Govender, V., Aveling, T.A.S., Kritzinger, Q., 2008. The effect of traditional storage methods on germination and vigour of maize (*Zea mays* L.) from northern KwaZulu-Natal and southern Mozambique. *S. Afr. J. Bot.* 74, 190–196. \ Dutta, S. K., Layek, J., Akoijam, R. S., Boopathi, T., Vanlalhmangaiha, Saha, S., & Prakash, N. (2019). Seaweed extract as natural priming agent for augmenting seed quality traits and yield in *Capsicum frutescens* L. *Journal of applied phycology*, 31, 3803-3813.

Examples

```
wt_of_seed_after_imbib <- 2.5  
initial_wt_of_seed <- 1.7  
seed_imbib(wt_of_seed_after_imbib, initial_wt_of_seed)
```

Index

* **Seed Imbibition Percentage**

SeedImbibition, [1](#)

seed_imbib (SeedImbibition), [1](#)

SeedImbibition, [1](#)