

Package ‘batsch’

July 22, 2025

Type Package

Title Real-Time PCR Data Sets by Batsch et al. (2008)

Version 0.1.1

Description Real-time quantitative polymerase chain reaction (qPCR) data sets by Batsch et al. (2008) <[doi:10.1186/1471-2105-9-95](https://doi.org/10.1186/1471-2105-9-95)>. This package provides five data sets, one for each PCR target: (i) rat SLC6A14, (ii) human SLC22A13, (iii) pig EMT, (iv) chicken ETT, and (v) human GAPDH. Each data set comprises a five-point, four-fold dilution series. For each concentration there are three replicates. Each amplification curve is 45 cycles long. Original raw data file: <https://static-content.springer.com/esm/art%3A10.1186%2F1471-2105-9-95/MediaObjects/12859_2007_2080_MOESM5_ESM.xls>.

License CC BY 4.0

Encoding UTF-8

LazyData true

RoxygenNote 7.3.1

Depends R (>= 2.10)

Imports tibble

URL <https://github.com/ramiromagno/batsch>, <https://rmagno.eu/batsch/>

BugReports <https://github.com/ramiromagno/batsch/issues>

NeedsCompilation no

Author Ramiro Magno [aut, cre] (ORCID: <<https://orcid.org/0000-0001-5226-3441>>),
Pattern Institute [cph, fnd]

Maintainer Ramiro Magno <rmagno@pattern.institute>

Repository CRAN

Date/Publication 2024-04-22 22:40:21 UTC

Contents

batsch	2
------------------	---

Index**4**

batsch

*qPCR data sets by Batsch et al. (2008)***Description**

Each data set comprises a five-point, four-fold dilution series. For each concentration there are three replicates. Each amplification curve is 45 cycles long.

A single reaction (total volume 10 µl) contained 1 µl master mix (LightCycler TaqMan Master; Roche 04735536001), 1 µmol/l each of forward and reverse primer, SYBR Green I at 1:30 dilution (Invitrogen S7563) or 50 nmol/l probe, and various amounts of cDNA or plasmid DNA. Contamination controls contained water instead of DNA. After enzyme activation (10 min, 95°C), thermocycling consisted of 45 cycles of 10 s at 95°C, 30 s at 55°C, and 1 s at 72°C; velocity of temperature change was 1.1°C/s. Please read the Methods section of Batsch et al. (2008) for more details.

Usage

SLC6A14r

SLC22A13h

EMTp

ETTch

GAPDHh

Format

Each data set is provided as a [tibble](#) with 675 rows and 9 variables:

plate Plate identifier. Because one plate was used per target, the name of the plate is the same as the values in **target**.

well Well identifier. Values are always NA (not available) for these data sets. This variable is kept nevertheless to be coherent with other data sets from other similar R data packages.

dye Either SYBR Green I master mix (Roche) ("SYBR") or TaqMan probe ("TaqMan").

target Target identifier: rat SLC6A14 ("SLC6A14r"), human SLC22A13 ("SLC22A13h"), pig EMT ("EMTp"), chicken ETT ("ETTch") or human GAPDH ("GAPDHh").

sample_type Sample type (all curves are standards, i.e. "std").

replicate Replicate identifier: 1 thru 3.

dilution Dilution factor. Higher number means greater dilution.

cycle PCR cycle.

fluor Raw fluorescence values.

Source

[doi:10.1186/14712105995](https://doi.org/10.1186/14712105995)

Examples

SLC6A14r

SLC22A13h

EMTp

ETTch

GAPDHh

Index

* **datasets**

batsch, [2](#)

batsch, [2](#)

EMTp (batsch), [2](#)

ETTch (batsch), [2](#)

GAPDHh (batsch), [2](#)

SLC22A13h (batsch), [2](#)

SLC6A14r (batsch), [2](#)

tibble, [2](#)