

Package ‘fpest’

July 22, 2025

Type Package

Title Estimating Finite Population Total

Version 0.1.1

Author S.Sampath

Maintainer S.Sampath <sampath1959@gmail.com>

Description Given the values of sampled units and selection probabilities
the `desraj` function in the package computes the estimated value of the total
as well as estimated variance.

License GPL-3

Encoding UTF-8

LazyData true

RoxygenNote 6.0.1

NeedsCompilation no

Repository CRAN

Date/Publication 2017-04-13 21:00:27 UTC

Contents

desraj	1
Index	3

desraj	<i>Desraj Ordered Estimator</i>
--------	---------------------------------

Description

Desraj estimator computes the estimated value of a finite population total when values of the study variable for the sampled units and the corresponding selection probabilities are given as per the order of selection. It is meant for use in Probability Proportional to Size Without Replacement Sampling Scheme.

Usage

```
desraj(y, p)
```

Arguments

y	vector of values of sampled units as per the order of selection
p	vector of selection probabilities as per the order of selection

References

Sampath,S.(2005) Sampling Theory and Methods,Alpha Science International, Ltd

Examples

```
y<-c(16,13,12,10)
p<-c(0.21, 0.34, 0.12,0.10)
desraj(y,p)
```

Index

desraj, [1](#)