

Package ‘llogistic’

July 22, 2025

Title The L-Logistic Distribution

Version 1.0.3

Description Density, distribution function, quantile function and random generation for the L-Logistic distribution with parameters m and ϕ . The parameter m is the median of the distribution.

Imports stats

Depends R ($\geq 3.3.0$)

License GPL-3

Encoding UTF-8

LazyData true

RoxygenNote 6.1.0

NeedsCompilation no

Author Rosineide Fernando da Paz [aut, cre],
Jorge Lu s Baz n [ctb]

Maintainer Rosineide Fernando da Paz <rfpaz2@gmail.com>

Repository CRAN

Date/Publication 2019-03-31 20:20:03 UTC

Contents

llogistic	1
Index	4

llogistic	<i>The L-Logistic Distribution</i>
-----------	------------------------------------

Description

Density, distribution function, quantile function and random generation for the L-Logistic distribution with parameters m and ϕ .

Usage

```
dllogistic(x, m, phi, log = FALSE)

pllogistic(q, m, phi, lower.tail = TRUE, log.p = FALSE)

qllogistic(p, m, phi, lower.tail = TRUE, log.p = FALSE)

rllogistic(n, m, phi)
```

Arguments

x, q	vector of quantiles.
m, phi	parameters of the L-Logistic distribution. The parameter m lies in the interval (0,1) and phi is positive.
log, log.p	logical; if TRUE, probabilities p are given as log(p).
lower.tail	logical; if TRUE (default), probabilities are $P[X \leq x]$, otherwise, $P[X > x]$.
p	vector of probabilities.
n	number of observations.

Details

The llogistic distribution has density

$$f(x) = \phi(1-m)^{\phi} m^{\phi} \phi (x(1-x))^{\phi-1} / ((1-m)^{\phi} x^{\phi} + m^{\phi} (1-x)^{\phi})^2,$$

for $0 < x < 1$, where m is a median of the distribution and phi is a shape parameter.

Value

dllogistic(x,m,phi) gives the density function, rllogistic(n,m,phi) gives n random variates and qllogistic(p,m,phi) gives the quantile.

Source

The L-Logistic distribution was introduced by Tadikamalla and Johnson (1982), which refer to this distribution as Logit-Logistic distribution. Here, we have a new parameterization of the Logit-Logistic with the median as a parameter.

References

Paz, R.F., Balakrishnan, N and Bazán, Jorge L. (2016). L-Logistic Distribution: Properties, Inference and an Application to Study Poverty and Inequality in Brazil. São Carlos: Universidade Federal de São Carlos. Technical-Scientific Report No. 261, Theory and Method. Sponsored by the Department of Statistical, <URL:<http://www.pipges.ufscar.br/publicacoes/relatorios-tecnicos/arquivos-1/rt261.pdf>>.

TADIKAMALLA, P. R.; JOHNSON, N. L. (1982). Systems of frequency curves generated by transformations of logistic variables. Biometrika, v. 69, n. 2, p. 461.

Examples

```
dllogistic(0.3, 0.5, 2)
pllogistic(0.7, 0.5, 2)
qllogistic(0.2, 0.5, 2)
rllogistic(10, 0.5, 2)
```

Index

`dllogistic(llogistic)`, [1](#)

`llogistic`, [1](#)

`pllogistic(llogistic)`, [1](#)

`qllogistic(llogistic)`, [1](#)

`rllogistic(llogistic)`, [1](#)