

Package ‘nlgm’

July 22, 2025

Type Package

Title Non Linear Growth Models

Version 1.0

Date 2022-11-25

Author Michail Tsagris [aut, cre],
Nikolaos Pandis [aut]

Maintainer Michail Tsagris <mtsagris@uoc.gr>

Depends R (>= 4.0)

Imports ggplot2, Rfast, Rfast2, stats

Description Six growth models are fitted using non-linear least squares. These are the Richards, the 3, 4 and 5 parameter logistic, the Gompertz and the Weibull growth models. Reference: Reddy T., Shkedy Z., van Rensburg C. J., Mwambi H., Debba P., Zuma K. and Manda, S. (2021). ``Short-term real-time prediction of total number of reported COVID-19 cases and deaths in South Africa: a data driven approach". BMC medical research methodology, 21(1), 1-11. <doi:10.1186/s12874-020-01165-x>.

License GPL (>= 2)

NeedsCompilation no

Repository CRAN

Date/Publication 2022-11-28 14:40:02 UTC

Contents

nlgm-package	2
Non-linear growth models	3
Parametric bootstrap prediction intervals	4
Plot of the fitted values of the non-linear growth models	6
Index	8

nlgm-package

This is an R package that fits 6 non linear growth models.

Description

Six growth models are fitted using non-linear least squares. The relevant reference is: Reddy T., Shkedy Z., van Rensburg C. J., Mwambi H., Debba P., Zuma K. and Manda, S. (2021). Short-term real-time prediction of total number of reported COVID-19 cases and deaths in South Africa: a data driven approach. BMC Medical Research methodology, 21(1): 1–11. <doi:10.1186/s12874-020-01165-x>.

Details

Package: nlgmcr Type: Package
Version: 1.0
Date: 2021-11-25
License: GPL-2

Maintainers

Michail Tsagris <mtsagris@uoc.gr>.

Note

Acknowledgments:

Author(s)

Michail Tsagris <mtsagris@uoc.gr> and Nikolaos Pandis <npandis@yahoo.com>.

References

Reddy T., Shkedy Z., van Rensburg C. J., Mwambi H., Debba P., Zuma K. and Manda, S. (2021). Short-term real-time prediction of total number of reported COVID-19 cases and deaths in South Africa: a data driven approach. BMC medical Research Methodology, 21(1): 1–11.

Non-linear growth models

Non-linear growth models

Description

Non-linear growth models.

Usage

```
nlgm(y, ti, type, ini, ti.ahead = NULL)
```

Arguments

<code>y</code>	A vector with the cumulative number of cases.
<code>ti</code>	A vector with the time, e.g. days.
<code>ini</code>	A vector with the initial values. The length varies according to the type of growth model.
<code>type</code>	The type of the growth model. "richards", "3logistic", "4logistic", "5logistic", "gomperz" or "weibull". See Reddy et al. (2021) for more information.
<code>ti.ahead</code>	If you want to make predictions for a number of days ahead, provide the number of days, a single number.

Details

Non-linear growth curves are fitted using least squares.

Value

A list including:

<code>mod</code>	The object as returned by the function <code>nls</code> .
<code>param</code>	A vector with the parameters of the chosen model.
<code>fit</code>	The fitted values.
<code>pred</code>	The future predictions if values for the argument "ti.ahead" were given, otherwise this is NULL.

Author(s)

Michail Tsagris and Nikolaos Pandis.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr> and Nikolaos Pandis <npandis@yahoo.com>.

References

Reddy T., Shkedy Z., van Rensburg C. J., Mwambi H., Debba P., Zuma K. and Manda, S. (2021). Short-term real-time prediction of total number of reported COVID-19 cases and deaths in South Africa: a data driven approach. BMC medical research methodology, 21(1), 1-11.

See Also

[boot.pred](#), [fit.plot](#)

Examples

```
## Data on Covid 19 for the first 96 days of Belgium
y <- c( 19, 38, 72, 125, 206, 316, 343, 407, 501, 600, 774, 1024, 1362, 1541, 1755, 2142,
        2564, 3098, 3811, 4473, 4942, 5428, 6756, 7951, 9150, 10513, 12031, 12875, 13558,
        15296, 16977, 18493, 19971, 21665, 22587, 23252, 25186, 26701, 28299, 30538, 32874,
        33903, 34427, 34964, 36524, 38157, 39831, 41225, 41947, 42390, 43666, 44936, 45713,
        46689, 47500, 47888, 48093, 48848, 49417, 49939, 50525, 50762, 51048, 51188, 51858,
        52404, 52956, 53398, 53881, 54121, 54239, 54715, 55110, 55431, 55736, 56082, 56229,
        56310, 56627, 56919, 57304, 57374, 57622, 57745, 57820, 58134, 58336, 58518, 58689,
        58854, 58915, 58964, 59023, 59204, 59363, 59535 )

ti <- 1:96
## Apply the 4-parameter logistic model
mod1 <- nlgm(y, ti, type = "4logistic", ini = c(60000, 1, 1, 35) )
## predict the cases 10 days ahead
mod2 <- nlgm(y, ti, type = "4logistic", ini = c(60000, 1, 1, 35), ti.ahead = 10 )
```

Parametric bootstrap prediction intervals

Parametric bootstrap prediction intervals

Description

Parametric bootstrap prediction intervals.

Usage

```
boot.pred(mod, type, ti, ti.ahead = 10, B = 1000, conf = 0.95, seed = NULL)
```

Arguments

<code>mod</code>	The "nls" object of the nlgm function.
<code>type</code>	The type of the growth model. "richards", "3logistic", "4logistic", "5logistic", "gompertz" or "weibull". See Reddy et al. (2021) for more information.
<code>ti</code>	A vector with the time, e.g. days.
<code>ti.ahead</code>	The future time points.

B	The number of bootstrap samples to draw. These samples are drawn from a Poisson distribution.
conf	The prediction level, set to 95% by default.
seed	Provide a seed number if you want, otherwise leave it NULL.

Details

Non-linear growth curves are fitted using least squares. Based on the model a parametric bootstrap is applied in order to construct prediction intervals. The fitted values act as the mean from which Poisson samples are drawn and the nlgm is fitted. Using this fitted model, the predicted number of cumulative cases, at the selected number of days ahead, are calculated. This process is repeated B times and in the end the prediction interval is computed by returning the tails of this bootstrap predicted values distribution.

Value

A list including:

est	A column with 3 columns, the mean of the bootstrap based predictions and the prediction interval.
pred	A matrix with B columns. Each column represents a bootstrap based set of prediction values.

Author(s)

Michail Tsagris and Nikolaos Pandis.

R implementation and documentation: Michail Tsagris <mtsagris@uoc.gr> and Nikolaos Pandis <npandis@yahoo.com>.

References

Reddy T., Shkedy Z., van Rensburg C. J., Mwambi H., Debba P., Zuma K. and Manda, S. (2021). Short-term real-time prediction of total number of reported COVID-19 cases and deaths in South Africa: a data driven approach. BMC medical research methodology, 21(1), 1-11.

See Also

[nlgm](#), [fit.plot](#)

Examples

```
## Data on the 96 first days of Belgium
y <- c( 19, 38, 72, 125, 206, 316, 343, 407, 501, 600, 774, 1024, 1362, 1541, 1755, 2142,
2564, 3098, 3811, 4473, 4942, 5428, 6756, 7951, 9150, 10513, 12031, 12875, 13558,
15296, 16977, 18493, 19971, 21665, 22587, 23252, 25186, 26701, 28299, 30538, 32874,
33903, 34427, 34964, 36524, 38157, 39831, 41225, 41947, 42390, 43666, 44936, 45713,
46689, 47500, 47888, 48093, 48848, 49417, 49939, 50525, 50762, 51048, 51188, 51858,
52404, 52956, 53398, 53881, 54121, 54239, 54715, 55110, 55431, 55736, 56082, 56229,
56310, 56627, 56919, 57304, 57374, 57622, 57745, 57820, 58134, 58336, 58518, 58689,
58854, 58915, 58964, 59023, 59204, 59363, 59535 )
```

```
ti <- 1:96
## Apply the 4-parameter logistic model
mod <- nlgm(y, ti, type = "4logistic", ini = c(60000, 1, 1, 35) )
preds <- boot.pred(mod, type = "4logistic", ti = ti, B = 100)
```

Plot of the fitted values of the non-linear growth models

Plot of the fitted values of the non-linear growth models

Description

Plot of the fitted values of the non-linear growth models.

Usage

```
fit.plot(mat)
```

Arguments

<code>mat</code>	A data.frame object with three columns, the observed cumulative cases, the time and the fitted values from a model. The names must be "cases", "ti" and "fit". See the examples below for an illustration.
------------------	--

Details

The scatter plot of the cumulative cases versus time is plotted. Then, lines with the fitted cases are added.

Value

A scatter plot of the cumulative cases versus time, including lines with the fitted cases.

Author(s)

Nikolaos Pandis and Michail Tsagris.

R implementation and documentation: Nikolaos Pandis <npandis@yahoo.com> and Michail Tsagris <mtsagris@uoc.gr>.

References

Reddy T., Shkedy Z., van Rensburg C. J., Mwambi H., Debba P., Zuma K. and Manda, S. (2021). Short-term real-time prediction of total number of reported COVID-19 cases and deaths in South Africa: a data driven approach. BMC medical research methodology, 21(1), 1-11.

See Also

[nlgm](#), [boot.pred](#)

Examples

```
## Data on the 96 first days of Belgium
y <- c( 19, 38, 72, 125, 206, 316, 343, 407, 501, 600, 774, 1024, 1362, 1541, 1755, 2142,
        2564, 3098, 3811, 4473, 4942, 5428, 6756, 7951, 9150, 10513, 12031, 12875, 13558,
        15296, 16977, 18493, 19971, 21665, 22587, 23252, 25186, 26701, 28299, 30538, 32874,
        33903, 34427, 34964, 36524, 38157, 39831, 41225, 41947, 42390, 43666, 44936, 45713,
        46689, 47500, 47888, 48093, 48848, 49417, 49939, 50525, 50762, 51048, 51188, 51858,
        52404, 52956, 53398, 53881, 54121, 54239, 54715, 55110, 55431, 55736, 56082, 56229,
        56310, 56627, 56919, 57304, 57374, 57622, 57745, 57820, 58134, 58336, 58518, 58689,
        58854, 58915, 58964, 59023, 59204, 59363, 59535 )
ti <- 1:96
mod <- nlgm(y, ti, type = "4logistic", ini = c(60000, 1, 1, 35) )
mat <- cbind(y, ti, mod$fit)
colnames(mat) <- c("cases", "ti", "fit")
mat <- as.data.frame(mat)
fit.plot(mat)
```

Index

* growth models

nlgm-package, 2

boot.pred, 4, 6

boot.pred (Parametric bootstrap
prediction intervals), 4

fit.plot, 4, 5

fit.plot (Plot of the fitted values of
the non-linear growth models),
6

nlgm, 4–6

nlgm (Non-linear growth models), 3

nlgm-package, 2

nls, 3

Non-linear growth models, 3

Parametric bootstrap prediction
intervals, 4

Plot of the fitted values of the
non-linear growth models, 6