

Package ‘spphpr’

July 23, 2025

Type Package

Title Spring Phenological Prediction

Version 1.1.5

Date 2025-06-20

Author Peijian Shi [aut, cre],
Zhenghong Chen [aut],
Jing Tan [aut],
Brady K. Quinn [aut]

Maintainer Peijian Shi <pjshi@njfu.edu.cn>

Description Predicts the occurrence times (in day-of-year) of spring phenological events. Three methods, including the accumulated degree days (ADD) method, the accumulated days transferred to a standardized temperature (ADTS) method, and the accumulated developmental progress (ADP) method, were used. See Shi et al. (2017a) <[doi:10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)> and Shi et al. (2017b) tails.

Depends R (>= 4.2.0)

License GPL (>= 2)

NeedsCompilation no

Repository CRAN

Date/Publication 2025-06-20 13:30:02 UTC

Contents

ADD	2
ADD2	5
ADD3	8
ADD4	11
ADP	14
ADP2	18
ADTS	22
ADTS2	25
apricotFFD	28
BJDAT	29

predADD	30
predADD2	32
predADP	34
predADP2	37
predADTS	41
predADTS2	44
spphpr	47
toDOY	49

Index	51
--------------	-----------

ADD	<i>Function for Implementing the Accumulated Degree Days Method Using Mean Daily Temperatures</i>
-----	---

Description

Estimates the starting date (S , in day-of-year) and base temperature (T_0 , in $^{\circ}\text{C}$) in the accumulated degree days method using mean daily air temperatures (Aono, 1993; Shi et al., 2017a, b).

Usage

```
ADD(S.pd = NULL, T0.arr, Year1, Time, Year2, DOY, Temp, DOY.ul = 120,
    fig.opt = TRUE, S.def = 54, verbose = TRUE)
```

Arguments

S.pd	the pre-determined starting date for thermal accumulation (in day-of-year)
T0.arr	the candidate base temperatures (in $^{\circ}\text{C}$)
Year1	the vector of the years in which a particular phenological event was recorded
Time	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
Year2	the vector of the years recording the climate data corresponding to the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Temp	the mean daily air temperature data (in $^{\circ}\text{C}$) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time
fig.opt	an optional argument to draw the figures associated with the determinations of the starting date and base temperature, and a comparison between the predicted and observed occurrence times
S.def	a mandatory definition of the starting date when (i) S.pd is NULL and (ii) the minimum correlation coefficient method fails to find a suitable starting date
verbose	an optional argument allowing users to suppress the printing of computation progress

Details

The default of `S.pd` is `NULL`. In this case, the date associated with the minimum correlation coefficient [between the mean of the mean daily temperatures (from a candidate starting date to the observed occurrence time) and the observed occurrence time] will be determined to be the starting date on the condition that it is smaller than the minimum phenological occurrence time. If the determined date associated with the minimum correlation coefficient is greater than the minimum phenological occurrence time, `S.def` will be used as the starting date. If `S.pd` is not `NULL`, the starting date will be directly set as `S.pd` irrespective of the minimum correlation coefficient method and the value of `S.def`. This means that `S.pd` is superior to `S.def` in determining the starting date.

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.ul` should be greater than or equal to the maximum `Time`.

Value

<code>S.arr</code>	the candidate starting dates (in day-of-year), whose default ranges from the minimum <code>DOY</code> to <code>min(DOY.ul, the maximum DOY)</code>
<code>cor.coef.arr</code>	the candidate correlation coefficients between the mean of the mean daily temperatures (from a candidate starting date to the observed occurrence time) and the observed occurrence time
<code>cor.coef</code>	the minimum correlation coefficient, i.e., <code>min(cor.coef.arr)</code>
<code>search.failure</code>	a value of 0 or 1 of showing whether the starting date is successfully determined by the minimum correlation coefficient method when <code>S.pd = NULL</code> , where 0 represents success and 1 represents failure
<code>mAADD.arr</code>	an vector saving the interannual mean of the annual accumulated degree days (AADD) values for each of the candidate base temperatures
<code>RMSE.arr</code>	a vector saving the candidate root-mean-square errors (in days) between the observed and predicted occurrence times for each of the candidate base temperatures
<code>AADD.arr</code>	the annual accumulated degree days (AADD) values in different years
<code>Year</code>	The overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time</code>	The observed occurrence times (day-of-year) in the overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time.pred</code>	the predicted occurrence times in different years
<code>S</code>	the determined starting date (day-of-year)
<code>T0</code>	the determined base temperature (in °C)
<code>AADD</code>	the expected annual accumulated degree days
<code>RMSE</code>	the smallest RMSE (in days) from the different candidate base temperatures
<code>unused.years</code>	the years that have phenological records but lack climate data

Note

The entire mean daily temperature data set for the spring of each year should be provided. AADD is represented by the mean of AADD.arr in the output. When the argument of S.pd is not NULL, the returned value of search.failure will be NA. When the argument of S.pd is NULL, and the minimum correlation coefficient method fails to find a suitable starting date, the argument of S.def is then defined as the determined starting date, i.e., the returned value of S. At the same time, the returned value of cor.coef is defined as NA.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).

Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001

Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063

See Also

[predADD](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
T0.arr0 <- seq(-5, 5, by = 0.1)
S.pd0 <- NULL

res1 <- ADD( S.pd = S.pd0, T0.arr = T0.arr0, Year1 = Year1.val, Time = Time.val,
            Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val,
            DOY.ul = DOY.ul.val, fig.opt = TRUE, S.def=54, verbose = TRUE )

res1
```

```

S0 <- res1$$.arr
r0 <- res1$cor.coef.arr

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( S0, r0, cex.lab = 1.5, cex.axis = 1.5, xlab = "Candidate starting date (day-of-year)",
      ylab="Correlation coefficient between the mean temperature and FFD", type="l" )
ind <- which.min(r0)
points(S0[ind], r0[ind], cex = 1.5, pch = 16)
text(S0[ind], r0[ind] + 0.1, bquote(paste(italic(S), " = ", .(S0[ind])), sep = "")), cex = 1.5)
par(par1)
par(par2)
par(par3)

resu1 <- ADD( S.pd = 65, T0.arr = seq(-10, 0, by = 0.1), Year1 = Year1.val, Time = Time.val,
             Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val,
             DOY.ul = DOY.ul.val, fig.opt = TRUE, S.def = 54, verbose = TRUE )

resu1

# graphics.off()

```

ADD2

Function for Implementing the Accumulated Degree Days Method Using Minimum and Maximum Daily Temperatures

Description

Estimates the starting date (S , in day-of-year) and base temperature (T_0 , in $^{\circ}\text{C}$) in the accumulated degree days method using minimum and maximum daily air temperatures (Aono, 1993; Shi et al., 2017a, b).

Usage

```
ADD2(S.pd = NULL, T0.arr, Year1, Time, Year2, DOY, Tmin, Tmax,
     DOY.ul = 120, fig.opt = TRUE, S.def = 54, verbose = TRUE)
```

Arguments

$S.pd$	the pre-determined starting date for thermal accumulation (in day-of-year)
$T0.arr$	the candidate base temperatures (in $^{\circ}\text{C}$)
$Year1$	the vector of the years in which a particular phenological event was recorded
$Time$	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years

Year2	the vector of the years recording the climate data corresponding to the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Tmin	the minimum daily air temperature data (in °C) corresponding to DOY
Tmax	the maximum daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time
fig.opt	an optional argument to draw the figures associated with the determinations of the starting date and base temperature, and a comparison between the predicted and observed occurrence times
S.def	a mandatory definition of the starting date when (i) S.pd is NULL and (ii) the minimum correlation coefficient method fails to find a suitable starting date
verbose	an optional argument allowing users to suppress the printing of computation progress

Details

The default of S.pd is NULL. In this case, the date associated with the minimum correlation coefficient [between the mean of the $(T_{min} + T_{max})/2$ values (from a candidate starting date to the observed occurrence time) and the observed occurrence time] will be determined to be the starting date on the condition that it is smaller than the minimum phenological occurrence time. If the determined date associated with the minimum correlation coefficient is greater than the minimum phenological occurrence time, S.def will be used as the starting date. If S.pd is not NULL, the starting date will be directly set as S.pd irrespective of the minimum correlation coefficient method and the value of S.def. This means that S.pd is superior to S.def in determining the starting date.

The function does not require that Year1 is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of DOY.ul should be greater than or equal to the maximum Time.

Value

S.arr	the candidate starting dates (in day-of-year), whose default ranges from the minimum DOY to $\min(\text{DOY.ul}, \text{the maximum DOY})$
cor.coef.arr	the candidate correlation coefficients between the mean of the $(T_{min} + T_{max})/2$ values (from a candidate starting date to the observed occurrence time) and the observed occurrence time
cor.coef	the minimum correlation coefficient, i.e., $\min(\text{cor.coef.arr})$
search.failure	a value of 0 or 1 of showing whether the starting date is successfully determined by the minimum correlation coefficient method when S.pd = NULL, where 0 represents success and 1 represents failure
mAADD.arr	an vector saving the interannual mean of the annual accumulated degree days (AADD) values for each of the candidate base temperatures
RMSE.arr	a vector saving the candidate root-mean-square errors (in days) between the observed and predicted occurrence times for each of the candidate base temperatures

AADD.arr	the annual accumulated degree days (AADD) values in different years
Year	The overlapping years between Year1 and Year2
Time	The observed occurrence times (day-of-year) in the overlapping years between Year1 and Year2
Time.pred	the predicted occurrence times in different years
S	the determined starting date (day-of-year)
T0	the determined base temperature (in °C)
AADD	the expected annual accumulated degree days
RMSE	the smallest RMSE (in days) from the different candidate base temperatures
unused.years	the years that have phenological records but lack climate data

Note

The entire minimum and maximum daily temperature data set for the spring of each year should be provided. AADD is represented by the mean of AADD.arr in the output. When the argument of S.pd is not NULL, the returned value of search.failure will be NA. When the argument of S.pd is NULL, and the minimum correlation coefficient method fails to find a suitable starting date, the argument of S.def is then defined as the determined starting date, i.e., the returned value of S. At the same time, the returned value of cor.coef is defined as NA.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063

See Also

[predADD2](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
```

```

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxDT
DOY.ul.val <- 120
T0.arr0 <- seq(3.5, 4, by = 0.1)
S.pd0 <- NULL

cand.res1 <- ADD2( S.pd = S.pd0, T0.arr = T0.arr0, Year1 = Year1.val, Time = Time.val,
                  Year2 = Year2.val, DOY = DOY.val, Tmin = Tmin.val, Tmax = Tmax.val,
                  DOY.ul = DOY.ul.val, fig.opt = TRUE, S.def=54, verbose = TRUE )

cand.res1

S0 <- cand.res1$S.arr
r0 <- cand.res1$cor.coef.arr

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( S0, r0, cex.lab = 1.5, cex.axis = 1.5, xlab = "Candidate starting date (day-of-year)",
      ylab="Correlation coefficient between the mean temperature and FFD", type="l" )
ind <- which.min(r0)
points(S0[ind], r0[ind], cex = 1.5, pch = 16)
text(S0[ind], r0[ind] + 0.1, bquote(paste(italic(S), " = ", .(S0[ind])), sep = "")), cex = 1.5)
par(par1)
par(par2)
par(par3)

# graphics.off()

```

ADD3

Function for Implementing the Accumulated Degree Days Method Using Mean Daily Temperatures for the Combinations of the Starting Date and Base Temperature

Description

Estimates the starting date (S , in day-of-year) and base temperature (T_0 , in $^{\circ}\text{C}$) in the accumulated degree days (ADD) method using mean daily air temperatures (Konno and Sugihara, 1986; Aono, 1993; Shi et al., 2017a, b).

Usage

```
ADD3( S.arr, T0.arr, Year1, Time, Year2, DOY, Temp, DOY.ul = 120,
      fig.opt = TRUE, verbose = TRUE )
```

Arguments

<code>S.arr</code>	the candidate starting dates for thermal accumulation (in day-of-year)
<code>T0.arr</code>	the candidate base temperatures (in °C)
<code>Year1</code>	the vector of the years in which a particular phenological event was recorded
<code>Time</code>	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
<code>Year2</code>	the vector of the years recording the climate data corresponding to the occurrence times
<code>DOY</code>	the vector of the dates (in day-of-year) for which climate data exist
<code>Temp</code>	the mean daily air temperature data (in °C) corresponding to DOY
<code>DOY.ul</code>	the upper limit of DOY used to predict the occurrence time
<code>fig.opt</code>	an optional argument to draw the figures associated with the determination of the combination the starting date and base temperature, and a comparison between the predicted and observed occurrence times
<code>verbose</code>	an optional argument allowing users to suppress the printing of computation progress

Details

When `fig.opt` is equal to `TRUE`, it will show the contours of the root-mean-square errors (RMSEs) based on different combinations of S and T_0 .

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.ul` should be greater than or equal to the maximum `Time`.

Value

<code>mAADD.mat</code>	a matrix consisting of the means of the annual accumulated degree days (AADD) values from the combinations of S and T_0
<code>RMSE.mat</code>	the matrix consisting of the RMSEs (in days) from different combinations of S and T_0
<code>AADD.arr</code>	the AADD values in different years associated with the smallest value in <code>RMSE.mat</code>
<code>Year</code>	The overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time</code>	The observed occurrence times (day-of-year) in the overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time.pred</code>	the predicted occurrence times in different years
<code>S</code>	the determined starting date (day-of-year)

T0	the determined base temperature (in °C)
AADD	the expected AADD
RMSE	the smallest RMSE (in days) in RMSE.mat from different combinations of S and T_0
unused.years	the years that have phenological records but lack climate data

Note

The entire mean daily temperature data set for the spring of each year should be provided. AADD is represented by the mean of AADD.arr in the output.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Konno, T., Sugihara, S. (1986) Temperature index for characterizing biological activity in soil and its application to decomposition of soil organic matter. *Bulletin of National Institute for Agro-Environmental Sciences* 1, 51–68 (in Japanese with English abstract).
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)

See Also

[predADD](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
S.arr0 <- seq(60, 70, by = 1)
```

```

T0.arr0 <- seq(-2, 5, by = 1)

RES1 <- ADD3( S.arr = S.arr0, T0.arr = T0.arr0, Year1 = Year1.val, Time = Time.val,
             Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val, DOY.ul = DOY.ul.val,
             fig.opt = TRUE, verbose = TRUE)

RES1

RMSE.mat0 <- RES1$RMSE.mat
RMSE.range <- range(RMSE.mat0)

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
image( S.arr0, T0.arr0, RMSE.mat0, col = terrain.colors(200), axes = TRUE,
       cex.axis = 1.5, cex.lab = 1.5, xlab = "Starting date (day-of-year)",
       ylab = expression(paste("Base temperature (", degree, "C)", sep = "")))
points( RES1$S, RES1$T0, cex = 1.5, pch = 16, col = 2 )
contour( S.arr0, T0.arr0, RMSE.mat0, levels = round(seq(RMSE.range[1],
              RMSE.range[2], len = 20), 4), add = TRUE, cex = 1.5, col = "#696969", labcex = 1.5)
par(par1)
par(par2)
par(par3)

# graphics.off()

```

ADD4

Function for Implementing the Accumulated Degree Days Method Using Minimum and Maximum Daily Temperatures for the Combinations of the Starting Date and Base Temperature

Description

Estimates the starting date (S , in day-of-year) and base temperature (T_0 , in $^{\circ}\text{C}$) in the accumulated degree days (ADD) method using minimum and maximum daily air temperatures (Konno and Sugihara, 1986; Aono, 1993; Shi et al., 2017a, b).

Usage

```
ADD4( S.arr, T0.arr, Year1, Time, Year2, DOY, Tmin, Tmax, DOY.ul = 120,
      fig.opt = TRUE, verbose = TRUE )
```

Arguments

$S.arr$	the candidate starting dates for thermal accumulation (in day-of-year)
$T0.arr$	the candidate base temperatures (in $^{\circ}\text{C}$)

Year1	the vector of the years in which a particular phenological event was recorded
Time	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
Year2	the vector of the years recording the climate data corresponding to the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Tmin	the minimum daily air temperature data (in °C) corresponding to DOY
Tmax	the maximum daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time
fig.opt	an optional argument to draw the figures associated with the determination of the combination the starting date and base temperature, and a comparison between the predicted and observed occurrence times
verbose	an optional argument allowing users to suppress the printing of computation progress

Details

When `fig.opt` is equal to `TRUE`, it will show the contours of the root-mean-square errors (RMSEs) based on different combinations of S and T_0 .

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.ul` should be greater than or equal to the maximum `Time`.

Value

<code>mAADD.mat</code>	a matrix consisting of the means of the annual accumulated degree days (AADD) values from the combinations of S and T_0
<code>RMSE.mat</code>	the matrix consisting of the RMSEs (in days) from different combinations of S and T_0
<code>AADD.arr</code>	the AADD values in different years associated with the smallest value in <code>RMSE.mat</code>
<code>Year</code>	The overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time</code>	The observed occurrence times (day-of-year) in the overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time.pred</code>	the predicted occurrence times in different years
<code>S</code>	the determined starting date (day-of-year)
<code>T0</code>	the determined base temperature (in °C)
<code>AADD</code>	the expected AADD
<code>RMSE</code>	the smallest RMSE (in days) in <code>RMSE.mat</code> from different combinations of S and T_0
<code>unused.years</code>	the years that have phenological records but lack climate data

Note

The entire mean daily temperature data set for the spring of each year should be provided. AADD is represented by the mean of AADD.arr in the output.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).

Konno, T., Sugihara, S. (1986) Temperature index for characterizing biological activity in soil and its application to decomposition of soil organic matter. *Bulletin of National Institute for Agro-Environmental Sciences* 1, 51–68 (in Japanese with English abstract).

Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)

Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)

See Also

[predADD2](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxDT
DOY.ul.val <- 120
S.arr0 <- seq(63, 66, by = 1)
T0.arr0 <- 3.8

RES2 <- ADD4( S.arr = S.arr0, T0.arr = T0.arr0, Year1 = Year1.val, Time = Time.val,
              Year2 = Year2.val, DOY = DOY.val, Tmin = Tmin.val, Tmax = Tmax.val,
              DOY.ul = DOY.ul.val, fig.opt = TRUE, verbose = TRUE)
```

RES2

```
RESU2 <- ADD4( S.arr = 65, T0.arr = seq(2, 6, by = 1), Year1 = Year1.val, Time = Time.val,
              Year2 = Year2.val, DOY = DOY.val, Tmin = Tmin.val, Tmax = Tmax.val,
              DOY.ul = DOY.ul.val, fig.opt = TRUE, verbose = TRUE)
```

RESU2

```
# graphics.off()
```

ADP

*Function for Implementing the Accumulated Developmental Progress
Method Using Mean Daily Temperatures*

Description

Estimates the starting date (S , in day-of-year) and the parameters of a developmental rate model in the accumulated developmental progress (ADP) method using mean daily air temperatures (Wagner et al., 1984; Shi et al., 2017a, b).

Usage

```
ADP( S.arr, expr, ini.val, Year1, Time, Year2, DOY, Temp, DOY.ul = 120,
     fig.opt = TRUE, control = list(), verbose = TRUE )
```

Arguments

<code>S.arr</code>	the candidate starting dates for thermal accumulation (in day-of-year)
<code>expr</code>	a user-defined model that is used in the accumulated developmental progress (ADP) method
<code>ini.val</code>	a vector or a list that saves the initial values of the parameters in <code>expr</code>
<code>Year1</code>	the vector of the years in which a particular phenological event was recorded
<code>Time</code>	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
<code>Year2</code>	the vector of the years recording the climate data corresponding to the occurrence times
<code>DOY</code>	the vector of the dates (in day-of-year) for which climate data exist
<code>Temp</code>	the mean daily air temperature data (in °C) corresponding to <code>DOY</code>
<code>DOY.ul</code>	the upper limit of <code>DOY</code> used to predict the occurrence time
<code>fig.opt</code>	an optional argument to draw the figures associated with the temperature-dependent developmental rate curve, the mean daily temperatures versus years, and a comparison between the predicted and observed occurrence times
<code>control</code>	the list of control parameters for using the <code>optim</code> function in the stats package
<code>verbose</code>	an optional argument allowing users to suppress the printing of computation progress

Details

It is better not to set too many candidate starting dates, as doing so will be time-consuming. If `expr` is selected as Arrhenius' equation, `S.arr` can be selected as the `S` obtained from the output of the `ADTS` function. Here, `expr` can be other nonlinear temperature-dependent developmental rate functions (see Shi et al. [2017b] for details). Further, `expr` can be any an arbitrary user-defined temperature-dependent developmental rate function, e.g., a function named `myfun`, but it needs to take the form of `myfun <- function(P, x){...}`, where `P` is the vector of the model parameter(s), and `x` is the vector of the predictor variable, i.e., the temperature variable.

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.ul` should be greater than or equal to the maximum `Time`.

Let r represent the temperature-dependent developmental rate, i.e., the reciprocal of the developmental duration required for completing a particular phenological event, at a constant temperature. In the accumulated developmental progress (ADP) method, when the annual accumulated developmental progress (AADP) reaches 100%, the phenological event is predicted to occur for each year. Let $AADP_i$ denote the AADP of the i th year, which equals

$$AADP_i = \sum_{j=S}^{E_i} r_{ij}(\mathbf{P}; T_{ij}),$$

where S represents the starting date (in day-of-year), E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year, \mathbf{P} is the vector of the model parameters in `expr`, and T_{ij} represents the mean daily temperature of the j th day of the i th year (in °C or K). In theory, $AADP_i = 100\%$, i.e., the AADP values of different years are a constant 100%. However, in practice, there is a certain deviation of $AADP_i$ from 100%. The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F r_{ij} = 100\%$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F r_{ij} < 100\%$ and $\sum_{j=S}^{F+1} r_{ij} > 100\%$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time. Let \hat{E}_i represent the predicted occurrence time of the i th year. Assume that there are n -year phenological records. When the starting date S and the temperature-dependent developmental rate model are known, the model parameters can be estimated using the Nelder-Mead optimization method (Nelder and Mead, 1965) to minimize the root-mean-square error (RMSE) between the observed and predicted occurrence times, i.e.,

$$\hat{\mathbf{P}} = \arg \min_{\mathbf{P}} \{\text{RMSE}\} = \arg \min_{\mathbf{P}} \sqrt{\frac{\sum_{i=1}^n (E_i - \hat{E}_i)^2}{n}}.$$

Because S is not determined, a group of candidate S values (in day-of-year) need to be provided. Assume that there are m candidate S values, i.e., $S_1, S_2, S_3, \dots, S_m$. For each S_q (where q ranges between 1 and m), we can obtain a vector of the estimated model parameters, $\hat{\mathbf{P}}_q$, by minimizing RMSE_q using the Nelder-Mead optimization method. Then we finally selected $\hat{\mathbf{P}}$ associated with $\min \{\text{RMSE}_1, \text{RMSE}_2, \text{RMSE}_3, \dots, \text{RMSE}_m\}$ as the target parameter vector.

Value

TDDR	the temperature-dependent developmental rate matrix consisting of the year, day-of-year, mean daily temperature and developmental rate columns
MAT	a matrix consisting of the candidate starting dates and the estimates of candidate model parameters with the corresponding RMSEs
Dev . accum	the calculated annual accumulated developmental progresses in different years
Year	The overlapping years between Year1 and Year2
Time	The observed occurrence times (day-of-year) in the overlapping years between Year1 and Year2
Time . pred	the predicted occurrence times in different years
S	the determined starting date (day-of-year)
par	the estimates of model parameters
RMSE	the RMSE (in days) between the observed and predicted occurrence times
unused . years	the years that have phenological records but lack climate data

Note

The entire mean daily temperature data set for the spring of each year should be provided. In TDDR, the first column of Year saves the years, the second column of DOY saves the day-of-year values, the third column of Temperature saves the mean daily air temperatures calculated between the starting date to the occurrence times, and the fourth column of Rate saves the calculated developmental rates corresponding to the mean daily temperatures.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Nelder, J.A., Mead, R. (1965) A simplex method for function minimization. *Computer Journal* 7, 308–313. doi:10.1093/comjnl/7.4.308
- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:10.1093/ee/12.2.482
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063
- Wagner, T.L., Wu, H.-I., Sharpe, P.J.H., Shcoolfield, R.M., Coulson, R.N. (1984) Modelling insect development rates: a literature review and application of a biophysical model. *Annals of the Entomological Society of America* 77, 208–225. doi:10.1093/aesa/77.2.208

See Also[predADP](#)**Examples**

```

data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
S.arr0 <- 47

#### Defines a re-parameterized Arrhenius' equation #####
Arrhenius.eqn <- function(P, x){
  B <- P[1]
  Ea <- P[2]
  R <- 1.987 * 10^(-3)
  x <- x + 273.15
  10^12*exp(B-Ea/(R*x))
}
#####

#### Provides the initial values of the parameter of Arrhenius' equation #####
ini.val0 <- list( B = 20, Ea = 14 )
#####

res5 <- ADP( S.arr = S.arr0, expr = Arrhenius.eqn, ini.val = ini.val0, Year1 = Year1.val,
            Time = Time.val, Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val,
            DOY.ul = DOY.ul.val, fig.opt = TRUE, control = list(trace = FALSE,
            reltol = 1e-12, maxit = 5000), verbose = TRUE )

res5

TDDR <- res5$TDDR
T <- TDDR$Temperature
r <- TDDR$Rate
Y <- res5$Year
DP <- res5$Dev.accum

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
Ind <- sort(T, index.return=TRUE)$ix
T1 <- T[Ind]
r1 <- r[Ind]

```

```

plot( T1, r1, cex.lab = 1.5, cex.axis = 1.5, pch = 1, cex = 1.5, col = 2, type = "l",
      xlab = expression(paste("Mean daily temperature (", degree, "C)", sep = "")),
      ylab = expression(paste("Calculated developmental rate (", {day}^{"-1"}, ") ", sep = "")) )
par(par1)
par(par2)
par(par3)

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( Y, DP * 100, xlab = "Year",
      ylab = "Accumulated developmental progress (%)",
      ylim = c(50, 150), cex.lab=1.5, cex.axis = 1.5, cex = 1.5 )
abline( h = 1 * 100, lwd = 1, col = 4, lty = 2 )
par(par1)
par(par2)
par(par3)

# graphics.off()

```

ADP2

*Function for Implementing the Accumulated Developmental Progress
Method Using Minimum and Maximum Daily Temperatures*

Description

Estimates the starting date (S , in day-of-year) and the parameters of a developmental rate model in the accumulated developmental progress (ADP) method using minimum and maximum daily air temperatures (Wagner et al., 1984; Shi et al., 2017a, b).

Usage

```
ADP2( S.arr, expr, ini.val, Year1, Time, Year2, DOY, Tmin, Tmax,
      DOY.ul = 120, fig.opt = TRUE, control = list(), verbose = TRUE )
```

Arguments

<code>S.arr</code>	the candidate starting dates for thermal accumulation (in day-of-year)
<code>expr</code>	a user-defined model that is used in the accumulated developmental progress (ADP) method
<code>ini.val</code>	a vector or a list that saves the initial values of the parameters in <code>expr</code>
<code>Year1</code>	the vector of the years in which a particular phenological event was recorded
<code>Time</code>	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
<code>Year2</code>	the vector of the years recording the climate data corresponding to the occurrence times

DOY	the vector of the dates (in day-of-year) for which climate data exist
Tmin	the minimum daily air temperature data (in °C) corresponding to DOY
Tmax	the maximum daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time
fig.opt	an optional argument to draw the figures associated with the temperature-dependent developmental rate curve, the mean daily temperatures versus years, and a comparison between the predicted and observed occurrence times
control	the list of control parameters for using the <code>optim</code> function in the stats package
verbose	an optional argument allowing users to suppress the printing of computation progress

Details

It is better not to set too many candidate starting dates, as doing so will be time-consuming. If `expr` is selected as Arrhenius' equation, `S.arr` can be selected as the `S` obtained from the output of the `ADTS2` function. Here, `expr` can be other nonlinear temperature-dependent developmental rate functions (see Shi et al. [2017b] for details). Further, `expr` can be any an arbitrary user-defined temperature-dependent developmental rate function, e.g., a function named `myfun`, but it needs to take the form of `myfun <- function(P, x){...}`, where `P` is the vector of the model parameter(s), and `x` is the vector of the predictor variable, i.e., the temperature variable.

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.ul` should be greater than or equal to the maximum Time.

Let r represent the temperature-dependent developmental rate, i.e., the reciprocal of the developmental duration required for completing a particular phenological event, at a constant temperature. In the accumulated developmental progress (ADP) method, when the annual accumulated developmental progress (AADP) reaches 100%, the phenological event is predicted to occur for each year. Let $AADP_i$ denote the AADP of the i th year, which equals

$$AADP_i = \sum_{j=S}^{E_i} \sum_{w=1}^{24} \frac{r_{ijw}(\mathbf{P}; T_{ijw})}{24},$$

where S represents the starting date (in day-of-year), E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year, \mathbf{P} is the vector of the model parameters in `expr`, T_{ijw} represents the estimated mean hourly temperature of the w th hour of the j th day of the i th year (in °C or K), and r_{ijw} represents the developmental rate (per hour) at T_{ijw} , which is transferred to r_{ij} (per day) by dividing 24. This packages takes the method proposed by Zohner et al. (2020) to estimate the mean hourly temperature for each of 24 hours:

$$T_w = \frac{T_{\max} - T_{\min}}{2} \sin\left(\frac{w\pi}{12} - \frac{\pi}{2}\right) + \frac{T_{\max} + T_{\min}}{2},$$

where w represents the w th hour of a day, and T_{\min} and T_{\max} represent the minimum and maximum temperatures of the day, respectively.

In theory, $AADP_i = 100\%$, i.e., the AADP values of different years are a constant 100%. However, in practice, there is a certain deviation of $AADP_i$ from 100%. The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F \sum_{w=1}^{24} r_{ijw}/24 = 100\%$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F \sum_{w=1}^{24} r_{ijw}/24 < 100\%$ and $\sum_{j=S}^{F+1} \sum_{w=1}^{24} r_{ijw}/24 > 100\%$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time. Let \hat{E}_i represent the predicted occurrence time of the i th year. Assume that there are n -year phenological records. When the starting date S and the temperature-dependent developmental rate model are known, the model parameters can be estimated using the Nelder-Mead optimization method (Nelder and Mead, 1965) to minimize the root-mean-square error (RMSE) between the observed and predicted occurrence times, i.e.,

$$\hat{\mathbf{P}} = \arg \min_{\mathbf{P}} \{\text{RMSE}\} = \arg \min_{\mathbf{P}} \sqrt{\frac{\sum_{i=1}^n (E_i - \hat{E}_i)^2}{n}}.$$

Because S is not determined, a group of candidate S values (in day-of-year) need to be provided. Assume that there are m candidate S values, i.e., $S_1, S_2, S_3, \dots, S_m$. For each S_q (where q ranges between 1 and m), we can obtain a vector of the estimated model parameters, $\hat{\mathbf{P}}_q$, by minimizing RMSE_q using the Nelder-Mead optimization method. Then we finally selected $\hat{\mathbf{P}}$ associated with $\min \{\text{RMSE}_1, \text{RMSE}_2, \text{RMSE}_3, \dots, \text{RMSE}_m\}$ as the target parameter vector.

Value

TDDR	the temperature-dependent developmental rate matrix consisting of the year, day-of-year, estimated mean daily temperature (= (Tmin + Tmax)/2) and developmental rate columns
MAT	a matrix consisting of the candidate starting dates and the estimates of candidate model parameters with the corresponding RMSEs
Dev.accum	the calculated annual accumulated developmental progresses in different years
Year	The overlapping years between Year1 and Year2
Time	The observed occurrence times (day-of-year) in the overlapping years between Year1 and Year2
Time.pred	the predicted occurrence times in different years
S	the determined starting date (day-of-year)
par	the estimates of model parameters
RMSE	the RMSE (in days) between the observed and predicted occurrence times
unused.years	the years that have phenological records but lack climate data

Note

The entire minimum and maximum daily temperature data set for the spring of each year should be provided. In TDDR, the first column of Year saves the years, the second column of DOY saves the day-of-year values, the third column of Temperature saves the estimated mean daily air temperatures (= (Tmin + Tmax)/2) from the starting date to the occurrence times, and the fourth column of Rate saves the calculated developmental rates corresponding to the estimated mean daily temperatures.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Nelder, J.A., Mead, R. (1965) A simplex method for function minimization. *Computer Journal* 7, 308–313. doi:[10.1093/comjnl/7.4.308](https://doi.org/10.1093/comjnl/7.4.308)
- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:[10.1093/ee/12.2.482](https://doi.org/10.1093/ee/12.2.482)
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)
- Wagner, T.L., Wu, H.-I., Sharpe, P.J.H., Shcoolfield, R.M., Coulson, R.N. (1984) Modelling insect development rates: a literature review and application of a biophysical model. *Annals of the Entomological Society of America* 77, 208–225. doi:[10.1093/aesa/77.2.208](https://doi.org/10.1093/aesa/77.2.208)
- Zohner, C.M., Mo, L., Sebal, V., Renner, S.S. (2020) Leaf-out in northern ecotypes of wide-ranging trees requires less spring warming, enhancing the risk of spring frost damage at cold limits. *Global Ecology and Biogeography* 29, 1056–1072. doi:[10.1111/geb.13088](https://doi.org/10.1111/geb.13088)

See Also

[predADP2](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxDT
DOY.ul.val <- 120
S.arr0 <- 46

#### Defines a re-parameterized Arrhenius' equation #####
Arrhenius.eqn <- function(P, x){
  B <- P[1]
  Ea <- P[2]
  R <- 1.987 * 10^(-3)
  x <- x + 273.15
```

```

    10^12*exp(B-Ea/(R*x))
  }
#####

#### Provides the initial values of the parameter of Arrhenius' equation ####
ini.val0 <- list( B = 20, Ea = 14 )
#####

# The usage is similar to that of the "ADP" function. There is only a need to
#   replace "Temp = Temp.val" with "Tmin = Tmin.val, Tmax = Tmax.val" when using
#   the "ADP2" function.

```

ADTS

Function for Implementing the Accumulated Days Transferred to a Standardized Temperature Method Using Mean Daily Temperatures

Description

Estimates the starting date (S , in day-of-year) and activation free energy (E_a , in $\text{kcal} \cdot \text{mol}^{-1}$) in the accumulated days transferred to a standardized temperature (ADTS) method using mean daily air temperatures (Konno and Sugihara, 1986; Aono, 1993; Shi et al., 2017a, b).

Usage

```
ADTS( S.arr, Ea.arr, Year1, Time, Year2, DOY, Temp, DOY.ul = 120,
      fig.opt = TRUE, verbose = TRUE )
```

Arguments

<code>S.arr</code>	the candidate starting dates for thermal accumulation (in day-of-year)
<code>Ea.arr</code>	the candidate activation free energy values (in $\text{kcal} \cdot \text{mol}^{-1}$)
<code>Year1</code>	the vector of the years in which a particular phenological event was recorded
<code>Time</code>	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
<code>Year2</code>	the vector of the years recording the climate data corresponding to the occurrence times
<code>DOY</code>	the vector of the dates (in day-of-year) for which climate data exist
<code>Temp</code>	the mean daily air temperature data (in $^{\circ}\text{C}$) corresponding to DOY
<code>DOY.ul</code>	the upper limit of DOY used to predict the occurrence time
<code>fig.opt</code>	an optional argument to draw the figures associated with the determination of the combination the starting date and activation free energy, and a comparison between the predicted and observed occurrence times
<code>verbose</code>	an optional argument allowing users to suppress the printing of computation progress

Details

When `fig.opt` is equal to `TRUE`, it will show the contours of the root-mean-square errors (RMSEs) based on different combinations of S and E_a .

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.u1` should be greater than or equal to the maximum `Time`.

Value

<code>mAADTS.mat</code>	a matrix consisting of the means of the annual accumulated days transferred to a standardized temperature (AADTS) values from the combinations of S and E_a
<code>RMSE.mat</code>	the matrix consisting of the RMSEs (in days) from different combinations of S and E_a
<code>AADTS.arr</code>	the AADTS values in different years associated with the smallest value in <code>RMSE.mat</code>
<code>Year</code>	The overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time</code>	The observed occurrence times (day-of-year) in the overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time.pred</code>	the predicted occurrence times in different years
<code>S</code>	the determined starting date (day-of-year)
<code>Ea</code>	the determined activation free energy value (in $\text{kcal}\cdot\text{mol}^{-1}$)
<code>AADD</code>	the expected AADTS
<code>RMSE</code>	the smallest RMSE (in days) in <code>RMSE.mat</code> from different combinations of S and E_a
<code>unused.years</code>	the years that have phenological records but lack climate data

Note

The entire mean daily temperature data set for the spring of each year should be provided. AADTS is represented by the mean of `AADTS.arr` in the output.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Konno, T., Sugihara, S. (1986) Temperature index for characterizing biological activity in soil and its application to decomposition of soil organic matter. *Bulletin of National Institute for Agro-Environmental Sciences* 1, 51–68 (in Japanese with English abstract).

Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)

Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)

See Also

[predADTS](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
S.arr0 <- seq(40, 60, by = 1)
Ea.arr0 <- seq(10, 20, by = 1)

res3 <- ADTS( S.arr = S.arr0, Ea.arr = Ea.arr0, Year1 = Year1.val, Time = Time.val,
              Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val, DOY.ul = DOY.ul.val,
              fig.opt = TRUE, verbose = TRUE)

res3

RMSE.mat0 <- res3$RMSE.mat
RMSE.range <- range(RMSE.mat0)

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
image( S.arr0, Ea.arr0, RMSE.mat0, col = terrain.colors(200), axes = TRUE,
       cex.axis = 1.5, cex.lab = 1.5, xlab = "Starting date (day-of-year)",
       ylab = expression(paste(italic(E["a"]), " (kcal" %.% "mol"^-1), ")"), sep = "")))
points( res3$S, res3$Ea, cex = 1.5, pch = 16, col = 2 )
contour( S.arr0, Ea.arr0, RMSE.mat0, levels = round(seq(RMSE.range[1],
              RMSE.range[2], len = 20), 4), add = TRUE, cex = 1.5, col = "#696969", labcex = 1.5)
par(par1)
par(par2)
par(par3)

resu3 <- ADTS( S.arr = 47, Ea.arr = seq(10, 20, by = 0.5), Year1 = Year1.val, Time = Time.val,
              Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val, DOY.ul = DOY.ul.val,
```



```

                                fig.opt = TRUE, verbose = TRUE)
resu3

# graphics.off()

```

ADTS2	<i>Function for Implementing the Accumulated Days Transferred to a Standardized Temperature Method Using Minimum and Maximum Daily Temperatures</i>
-------	---

Description

Estimates the starting date (S , in day-of-year) and activation free energy (E_a , in $\text{kcal} \cdot \text{mol}^{-1}$) in the accumulated days transferred to a standardized temperature (ADTS) method using minimum and maximum daily air temperatures (Konno and Sugihara, 1986; Aono, 1993; Shi et al., 2017a, b).

Usage

```
ADTS2( S.arr, Ea.arr, Year1, Time, Year2, DOY, Tmin, Tmax,
       DOY.ul = 120, fig.opt = TRUE, verbose = TRUE )
```

Arguments

<code>S.arr</code>	the candidate starting dates for thermal accumulation (in day-of-year)
<code>Ea.arr</code>	the candidate activation free energy values (in $\text{kcal} \cdot \text{mol}^{-1}$)
<code>Year1</code>	the vector of the years in which a particular phenological event was recorded
<code>Time</code>	the vector of the occurrence times (in day-of-year) of a particular phenological event across many years
<code>Year2</code>	the vector of the years recording the climate data corresponding to the occurrence times
<code>DOY</code>	the vector of the dates (in day-of-year) for which climate data exist
<code>Tmin</code>	the minimum daily air temperature data (in $^{\circ}\text{C}$) corresponding to DOY
<code>Tmax</code>	the maximum daily air temperature data (in $^{\circ}\text{C}$) corresponding to DOY
<code>DOY.ul</code>	the upper limit of DOY used to predict the occurrence time
<code>fig.opt</code>	an optional argument to draw the figures associated with the determination of the combination the starting date and activation free energy, and a comparison between the predicted and observed occurrence times
<code>verbose</code>	an optional argument allowing users to suppress the printing of computation progress

Details

When `fig.opt` is equal to `TRUE`, it will show the contours of the root-mean-square errors (RMSEs) based on different combinations of S and E_a .

The function does not require that `Year1` is the same as `unique(Year2)`, and the intersection of the two vectors of years will be kept. The unused years that have phenological records but lack climate data will be showed in `unused.years` in the returned list.

The numerical value of `DOY.u1` should be greater than or equal to the maximum `Time`.

Value

<code>mAADTS.mat</code>	a matrix consisting of the means of the annual accumulated days transferred to a standardized temperature (AADTS) values from the combinations of S and E_a
<code>RMSE.mat</code>	the matrix consisting of the RMSEs (in days) from different combinations of S and E_a
<code>AADTS.arr</code>	the AADTS values in different years associated with the smallest value in <code>RMSE.mat</code>
<code>Year</code>	The overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time</code>	The observed occurrence times (day-of-year) in the overlapping years between <code>Year1</code> and <code>Year2</code>
<code>Time.pred</code>	the predicted occurrence times in different years
<code>S</code>	the determined starting date (day-of-year)
<code>Ea</code>	the determined activation free energy value (in $\text{kcal}\cdot\text{mol}^{-1}$)
<code>AADD</code>	the expected AADTS
<code>RMSE</code>	the smallest RMSE (in days) in <code>RMSE.mat</code> from different combinations of S and E_a
<code>unused.years</code>	the years that have phenological records but lack climate data

Note

The entire minimum and maximum daily temperature data set for the spring of each year should be provided. AADTS is represented by the mean of `AADTS.arr` in the output.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Konno, T., Sugihara, S. (1986) Temperature index for characterizing biological activity in soil and its application to decomposition of soil organic matter. *Bulletin of National Institute for Agro-Environmental Sciences* 1, 51–68 (in Japanese with English abstract).

Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)

Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)

See Also

[predADTS2](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT

Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxDT
DOY.ul.val <- 120
S.arr0 <- seq(45, 47, by = 1)
Ea.arr0 <- seq(20, 24, by = 0.5)

cand.res3 <- ADTS2( S.arr = S.arr0, Ea.arr = Ea.arr0, Year1 = Year1.val, Time = Time.val,
  Year2 = Year2.val, DOY = DOY.val, Tmin = Tmin.val, Tmax = Tmax.val,
  DOY.ul = DOY.ul.val, fig.opt = TRUE, verbose = TRUE)

cand.res3

RMSE.mat0 <- cand.res3$RMSE.mat
RMSE.range <- range(RMSE.mat0)

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
image( S.arr0, Ea.arr0, RMSE.mat0, col = terrain.colors(200), axes = TRUE,
  cex.axis = 1.5, cex.lab = 1.5, xlab = "Starting date (day-of-year)",
  ylab = expression(paste(italic(E["a"]), " (kcal" %.% "mol"^-1), ") ", sep = "")))
points( cand.res3$S, cand.res3$Ea, cex = 1.5, pch = 16, col = 2 )
contour( S.arr0, Ea.arr0, RMSE.mat0, levels = round(seq(RMSE.range[1],
  RMSE.range[2], len = 20), 4), add = TRUE, cex = 1.5, col = "#696969", labcex = 1.5)
par(par1)
par(par2)
par(par3)

# graphics.off()
```

apricotFFD

First flowering date records of Prunus armeniaca

Description

The data consist of the first flowering date records of *Prunus armeniaca* at the Summer Palace (39°54'38" N, 116°8'28" E, 50 m a.s.l.) in Beijing, China between 1963 and 2010 with the exception of 1969–1971, and 1997–2002. **Data source:** Chinese Phenological Observation Network (Guo et al., 2015).

Usage

```
data(apricotFFD)
```

Details

In the data set, there are two columns of vectors: Year and Time. Year saves the recording years; and Time saves the 1963–2010 first flowering dates of *Prunus armeniaca* (in day-of-year).

References

Guo, L., Xu, J., Dai, J., Cheng, J., Wu, H., Luedeling, E. (2015) Statistical identification of chilling and heat requirements for apricotflower buds in Beijing, China. *Scientia Horticulturae* 195, 138–144. doi:[10.1016/j.scienta.2015.09.006](https://doi.org/10.1016/j.scienta.2015.09.006)

Examples

```
data(apricotFFD)
attach(apricotFFD)

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( Year, Time, asp = 1, cex.lab = 1.5, cex.axis = 1.5,
      xlab = "Year", ylab = "First flowering date (day-of-year)" )
par(par1)
par(par2)
par(par3)

# graphics.off()
```

BJDAT

Daily Air Temperature Data of Beijing from 1952 to 2012.

Description

The data include the mean, minimum, and maximum daily temperatures (in °C) of Beijing between 1952 and 2012. **Data source:** China Meteorological Data Service Centre (<https://data.cma.cn/en>).

Usage

```
data(BJDAT)
```

Details

In the data set, there are seven columns of vectors: Year, Month, Day, DOY, MDT, MinDT, and MaxDT. Year saves the recording years; Month saves the recording months; Day saves the recording days; DOY saves the dates in day-of-year; MDT saves the mean daily temperatures (in °C) corresponding to DOY; MinDT saves the minimum daily temperatures (in °C) corresponding to DOY; MaxDT saves the maximum daily temperatures (in °C) corresponding to DOY.

References

Guo, L., Xu, J., Dai, J., Cheng, J., Wu, H., Luedeling, E. (2015) Statistical identification of chilling and heat requirements for apricotflower buds in Beijing, China. *Scientia Horticulturae* 195, 138–144. doi:[10.1016/j.scienta.2015.09.006](https://doi.org/10.1016/j.scienta.2015.09.006)

Examples

```
data(BJDAT)
attach(BJDAT)

x <- as.numeric( tapply(DOY, DOY, mean) )
y <- as.numeric( tapply(MDT, DOY, mean) )
y.sd <- as.numeric( tapply(MDT, DOY, sd) )

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( x, y, cex = 1.5, xlim = c(0, 367), ylim = c(-10, 30),
      cex.lab = 1.5, cex.axis = 1.5, type = "n", xlab = "Day-of-year",
      ylab = expression(paste("Mean daily temperature (", degree, "C)", sep="")) )
for(i in 1:length(x)){
  lines(c(x[i], x[i]), c(y[i]-y.sd[i], y[i]+y.sd[i]), col=4)
}
points(x, y, cex = 1.5)
par(par1)
par(par2)
```

```
par(par3)

# graphics.off()
```

predADD	<i>Prediction Function of the Accumulated Degree Days Method Using Mean Daily Temperatures</i>
---------	--

Description

Predicts the occurrence times using the accumulated degree days method based on observed or predicted mean daily air temperatures (Aono, 1993; Shi et al., 2017a, b).

Usage

```
predADD(S, T0, AADD, Year2, DOY, Temp, DOY.ul = 120)
```

Arguments

S	the starting date for thermal accumulation (in day-of-year)
T0	the base temperature (in °C)
AADD	the expected annual accumulated degree days
Year2	the vector of the years recording the climate data for predicting the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Temp	the mean daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time

Details

In the accumulated degree days (ADD) method (Shi et al., 2017a, b), the starting date (S), the base temperature (T_0), and the annual accumulated degree days (AADD, which is denoted by k) are assumed to be constants across different years. Let k_i denote the AADD of the i th year, which equals

$$k_i = \sum_{j=S}^{E_i} (T_{ij} - T_0),$$

where E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year, and T_{ij} represents the mean daily temperature of the j th day of the i th year (in °C). If $T_{ij} \leq T_0$, $T_{ij} - T_0$ is defined to be zero. In theory, $k_i = k$, i.e., the AADD values of different years are a constant. However, in practice, there is a certain deviation of k_i from k that is estimated by \bar{k} (i.e., the mean of the k_i values). The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F (T_{ij} - T_0) = \bar{k}$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F (T_{ij} - T_0) < \bar{k}$ and $\sum_{j=S}^{F+1} (T_{ij} - T_0) > \bar{k}$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time.

Value

Year the years with climate data
 Time.pred the predicted occurrence times (day-of-year) in different years

Note

The entire mean daily temperature data set for the spring of each year should be provided.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:10.1093/ee/12.2.482
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063

See Also

[ADD](#), [ADD3](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
S.val <- 65
T0.val <- -0.5
AADD.val <- 235.5282

res2 <- predADD( S = S.val, T0 = T0.val, AADD = AADD.val,
  Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val,
```

```

DOY.ul = DOY.ul.val )
res2

ind1 <- res2$Year %in% intersect(res2$Year, Year1.val)
ind2 <- Year1.val %in% intersect(res2$Year, Year1.val)
RMSE1 <- sqrt( sum((Time.val[ind2]-res2$Time.pred[ind1])^2) / length(Time.val[ind2]) )
RMSE1

```

predADD2	<i>Prediction Function of the Accumulated Degree Days Method Using Minimum and Maximum Daily Temperatures</i>
----------	---

Description

Predicts the occurrence times using the accumulated degree days method based on observed or predicted minimum and maximum daily air temperatures (Aono, 1993; Shi et al., 2017a, b).

Usage

```
predADD2(S, T0, AADD, Year2, DOY, Tmin, Tmax, DOY.ul = 120)
```

Arguments

S	the starting date for thermal accumulation (in day-of-year)
T0	the base temperature (in °C)
AADD	the expected annual accumulated degree days
Year2	the vector of the years recording the climate data for predicting the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Tmin	the minimum daily air temperature data (in °C) corresponding to DOY
Tmax	the maximum daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time

Details

In the accumulated degree days (ADD) method (Shi et al., 2017a, b), the starting date (S), the base temperature (T_0), and the annual accumulated degree days (AADD, which is denoted by k) are assumed to be constants across different years. Let k_i denote the AADD of the i th year, which equals

$$k_i = \sum_{j=S}^{E_i} \sum_{w=1}^{24} \frac{(T_{ijw} - T_0)}{24},$$

where E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year, and T_{ijw} represents the estimated mean hourly temperature of the w th hour of the j th day of the i th year (in °C). If $T_{ijw} \leq T_0$, $T_{ijw} - T_0$ is defined to be zero. This package takes the method proposed by Zohner et al. (2020) to estimate the mean hourly temperature (T_w) for each of 24 hours:

$$T_w = \frac{T_{\max} - T_{\min}}{2} \sin\left(\frac{w\pi}{12} - \frac{\pi}{2}\right) + \frac{T_{\max} + T_{\min}}{2},$$

where w represents the w th hour of a day, and T_{\min} and T_{\max} represent the minimum and maximum temperatures of the day, respectively.

In theory, $k_i = k$, i.e., the AADD values of different years are a constant. However, in practice, there is a certain deviation of k_i from k that is estimated by \bar{k} (i.e., the mean of the k_i values). The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F \sum_{w=1}^{24} (T_{ijw} - T_0) / 24 = \bar{k}$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F \sum_{w=1}^{24} (T_{ijw} - T_0) / 24 < \bar{k}$ and $\sum_{j=S}^{F+1} \sum_{w=1}^{24} (T_{ijw} - T_0) / 24 > \bar{k}$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time.

Value

Year	the years with climate data
Time.pred	the predicted occurrence times (day-of-year) in different years

Note

The entire minimum and maximum daily temperature data set for the spring of each year should be provided.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjmwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:10.1093/ee/12.2.482
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063
- Zohner, C.M., Mo, L., Sebal, V., Renner, S.S. (2020) Leaf-out in northern ecotypes of wide-ranging trees requires less spring warming, enhancing the risk of spring frost damage at cold limits. *Global Ecology and Biogeography* 29, 1056–1072. doi:10.1111/geb.13088

See Also[ADD2](#), [ADD4](#)**Examples**

```

data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxDT
DOY.ul.val <- 120
S.val <- 65
T0.val <- 3.8
AADD.val <- 136.5805

cand.res2 <- predADD2( S = S.val, T0 = T0.val, AADD = AADD.val, Year2 = Year2.val,
                      DOY = DOY.val, Tmin = Tmin.val, Tmax = Tmax.val,
                      DOY.ul = DOY.ul.val )

cand.res2

ind1 <- cand.res2$Year %in% intersect(cand.res2$Year, Year1.val)
ind2 <- Year1.val %in% intersect(cand.res2$Year, Year1.val)
RMSE1 <- sqrt( sum((Time.val[ind2]-cand.res2$Time.pred[ind1])^2) / length(Time.val[ind2]) )
RMSE1

```

predADP

*Prediction Function of the Accumulated Developmental Progress
Method Using Mean Daily Temperatures*

Description

Predicts the occurrence times using the accumulated developmental progress (ADP) method based on observed or predicted mean daily air temperatures (Wagner et al., 1984; Shi et al., 2017a, b).

Usage

```
predADP(S, expr, theta, Year2, DOY, Temp, DOY.ul = 120)
```

Arguments

S	the starting date for thermal accumulation (in day-of-year)
expr	a user-defined model that is used in the accumulated developmental progress (ADP) method
theta	a vector saves the numerical values of the parameters in expr
Year2	the vector of the years recording the climate data for predicting the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Temp	the mean daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time

Details

Organisms exhibiting phenological events in early spring often experience several cold days during their development. In this case, Arrhenius' equation (Shi et al., 2017a, b, and references therein) has been recommended to describe the effect of the absolute temperature (T in Kelvin [K]) on the developmental rate (r):

$$r = \exp \left(B - \frac{E_a}{RT} \right),$$

where E_a represents the activation free energy (in $\text{kcal} \cdot \text{mol}^{-1}$); R is the universal gas constant ($= 1.987 \text{ cal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$); B is a constant. To maintain consistency between the units used for E_a and R , we need to re-assign R to be 1.987×10^{-3} , making its unit $1.987 \times 10^{-3} \text{ kcal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$ in the above formula.

In the accumulated developmental progress (ADP) method, when the annual accumulated developmental progress (AADP) reaches 100%, the phenological event is predicted to occur for each year. Let AADP_i denote the AADP of the i th year, which equals

$$\text{AADP}_i = \sum_{j=S}^{E_i} r_{ij},$$

where E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year. If the temperature-dependent developmental rate follows Arrhenius' equation, the AADP of the i th year is equal to

$$\text{AADP}_i = \sum_{j=S}^{E_i} \exp \left(B - \frac{E_a}{RT_{ij}} \right),$$

where T_{ij} represents the mean daily temperature of the j th day of the i th year (in K). In theory, $\text{AADP}_i = 100\%$, i.e., the AADP values of different years are a constant 100%. However, in practice, there is a certain deviation of AADP_i from 100%. The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F r_{ij} = 100\%$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F r_{ij} < 100\%$ and $\sum_{j=S}^{F+1} r_{ij} > 100\%$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time.

The argument of `expr` can be any an arbitrary user-defined temperature-dependent developmental rate function, e.g., a function named `myfun`, but it needs to take the form of `myfun <- function(P, x){...}`, where `P` is the vector of the model parameter(s), and `x` is the vector of the predictor variable, i.e., the temperature variable.

Value

Year	the years with climate data
Time.pred	the predicted occurrence times (day-of-year) in different years

Note

The entire mean daily temperature data set for the spring of each year should be provided. It should be noted that the unit of Temp in **Arguments** is °C, not K. In addition, when using Arrhenius' equation to describe r , to reduce the size of B in this equation, Arrhenius' equation is multiplied by 10^{12} in calculating the AADP value for each year, i.e.,

$$\text{AADP}_i = \sum_{j=S}^{E_i} \left[10^{12} \cdot \exp \left(B - \frac{E_a}{RT_{ij}} \right) \right].$$

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:[10.1093/ee/12.2.482](https://doi.org/10.1093/ee/12.2.482)
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)
- Wagner, T.L., Wu, H.-I., Sharpe, P.J.H., Shcoolfield, R.M., Coulson, R.N. (1984) Modelling insect developmental rates: a literature review and application of a biophysical model. *Annals of the Entomological Society of America* 77, 208–225. doi:[10.1093/aesa/77.2.208](https://doi.org/10.1093/aesa/77.2.208)

See Also

[ADP](#)

Examples

```

data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
S.val <- 47

# Defines a re-parameterized Arrhenius' equation
Arrhenius.eqn <- function(P, x){
  B <- P[1]
  Ea <- P[2]
  R <- 1.987 * 10^(-3)
  x <- x + 273.15
  10^12*exp(B-Ea/(R*x))
}

P0 <- c(-4.3787, 15.0431)
T2 <- seq(-10, 20, len = 2000)
r2 <- Arrhenius.eqn(P = P0, x = T2)

dev.new()
par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( T2, r2, cex.lab = 1.5, cex.axis = 1.5, pch = 1, cex = 1.5, col = 2, type = "l",
      xlab = expression(paste("Temperature (", degree, "C)", sep = "")),
      ylab = expression(paste("Developmental rate (", {day}^{"-1"}, ") ", sep = "")) )
par(par1)
par(par2)
par(par3)

res6 <- predADP( S = S.val, expr = Arrhenius.eqn, theta = P0, Year2 = Year2.val,
                DOY = DOY.val, Temp = Temp.val, DOY.ul = DOY.ul.val )
res6

ind5 <- res6$Year %in% intersect(res6$Year, Year1.val)
ind6 <- Year1.val %in% intersect(res6$Year, Year1.val)
RMSE3 <- sqrt( sum((Time.val[ind6]-res6$Time.pred[ind5])^2) / length(Time.val[ind6]) )
RMSE3

```

Description

Predicts the occurrence times using the accumulated developmental progress (ADP) method based on observed or predicted minimum and maximum daily air temperatures (Wagner et al., 1984; Shi et al., 2017a, b).

Usage

```
predADP2(S, expr, theta, Year2, DOY, Tmin, Tmax, DOY.ul = 120)
```

Arguments

S	the starting date for thermal accumulation (in day-of-year)
expr	a user-defined model that is used in the accumulated developmental progress (ADP) method
theta	a vector saves the numerical values of the parameters in expr
Year2	the vector of the years recording the climate data for predicting the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Tmin	the minimum daily air temperature data (in °C) corresponding to DOY
Tmax	the maximum daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time

Details

Organisms exhibiting phenological events in early spring often experience several cold days during their development. In this case, Arrhenius' equation (Shi et al., 2017a, b, and references therein) has been recommended to describe the effect of the absolute temperature (T in Kelvin [K]) on the developmental rate (r):

$$r = \exp \left(B - \frac{E_a}{RT} \right),$$

where E_a represents the activation free energy (in $\text{kcal} \cdot \text{mol}^{-1}$); R is the universal gas constant ($= 1.987 \text{ cal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$); B is a constant. To maintain consistency between the units used for E_a and R , we need to re-assign R to be 1.987×10^{-3} , making its unit $1.987 \times 10^{-3} \text{ kcal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$ in the above formula.

In the accumulated developmental progress (ADP) method, when the annual accumulated developmental progress (AADP) reaches 100%, the phenological event is predicted to occur for each year. Let AADP_i denote the AADP of the i th year, which equals

$$\text{AADP}_i = \sum_{j=S}^{E_i} \sum_{w=1}^{24} \frac{r_{ijw}}{24},$$

where E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year. r_{ijw} is the developmental rate (per hour), which is transferred to r_{ij} (per day) by dividing 24. If the temperature-dependent developmental rate follows Arrhenius' equation, the AADP of the i th year is equal to

$$\text{AADP}_i = \sum_{j=S}^{E_i} \sum_{w=1}^{24} \left\{ \frac{1}{24} \exp \left(B - \frac{E_a}{R T_{ijw}} \right) \right\},$$

where T_{ijw} represents the estimated mean hourly temperature of the w th hour of the j th day of the i th year (in K). This packages takes the method proposed by Zohner et al. (2020) to estimate the mean hourly temperature (T_w) for each of 24 hours:

$$T_w = \frac{T_{\max} - T_{\min}}{2} \sin \left(\frac{w\pi}{12} - \frac{\pi}{2} \right) + \frac{T_{\max} + T_{\min}}{2},$$

where w represents the w th hour of a day, and T_{\min} and T_{\max} represent the minimum and maximum temperatures of the day, respectively.

In theory, $\text{AADP}_i = 100\%$, i.e., the AADP values of different years are a constant 100%. However, in practice, there is a certain deviation of AADP_i from 100%. The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F \sum_{w=1}^{24} (r_{ijw}/24) = 100\%$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F \sum_{w=1}^{24} (r_{ijw}/24) < 100\%$ and $\sum_{j=S}^{F+1} \sum_{w=1}^{24} (r_{ijw}/24) > 100\%$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time.

The argument of `expr` can be any an arbitrary user-defined temperature-dependent developmental rate function, e.g., a function named `myfun`, but it needs to take the form of `myfun <- function(P, x){...}`, where `P` is the vector of the model parameter(s), and `x` is the vector of the predictor variable, i.e., the temperature variable.

Value

Year	the years with climate data
Time.pred	the predicted occurrence times (day-of-year) in different years

Note

The entire minimum and maximum daily temperature data set for the spring of each year should be provided. It should be noted that the unit of T_{\min} and T_{\max} in **Arguments** is $^{\circ}\text{C}$, not K. In addition, when using Arrhenius' equation to describe r , to reduce the size of B in this equation, Arrhenius' equation is multiplied by 10^{12} in calculating the AADP value for each year, i.e.,

$$\text{AADP}_i = \sum_{j=S}^{E_i} \sum_{w=1}^{24} \left[10^{12} \cdot \frac{1}{24} \cdot \exp \left(B - \frac{E_a}{R T_{ijw}} \right) \right].$$

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:[10.1093/ee/12.2.482](https://doi.org/10.1093/ee/12.2.482)
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)
- Wagner, T.L., Wu, H.-I., Sharpe, P.J.H., Shcoolfield, R.M., Coulson, R.N. (1984) Modelling insect development rates: a literature review and application of a biophysical model. *Annals of the Entomological Society of America* 77, 208–225. doi:[10.1093/aesa/77.2.208](https://doi.org/10.1093/aesa/77.2.208)
- Zohner, C.M., Mo, L., Sebal, V., Renner, S.S. (2020) Leaf-out in northern ecotypes of wide-ranging trees requires less spring warming, enhancing the risk of spring frost damage at cold limits. *Global Ecology and Biogeography* 29, 1056–1072. doi:[10.1111/geb.13088](https://doi.org/10.1111/geb.13088)

See Also

[ADP2](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxDT
DOY.ul.val <- 120
S.val <- 46

# Defines a re-parameterized Arrhenius' equation
Arrhenius.eqn <- function(P, x){
  B <- P[1]
  Ea <- P[2]
  R <- 1.987 * 10^(-3)
  x <- x + 273.15
  10^12*exp(B-Ea/(R*x))
}

P0 <- c(8.220327, 22.185942)
T2 <- seq(-10, 20, len = 2000)
r2 <- Arrhenius.eqn(P = P0, x = T2)

dev.new()
```



```

par1 <- par(family="serif")
par2 <- par(mar=c(5, 5, 2, 2))
par3 <- par(mgp=c(3, 1, 0))
plot( T2, r2, cex.lab = 1.5, cex.axis = 1.5, pch = 1, cex = 1.5, col = 2, type = "l",
      xlab = expression(paste("Temperature (", degree, "C)", sep = "")),
      ylab = expression(paste("Developmental rate (", {day}^{"-1"}, ")", sep="")) )
par(par1)
par(par2)
par(par3)

cand.res6 <- predADP2( S = S.val, expr = Arrhenius.eqn, theta = P0, Year2 = Year2.val,
                      DOY = DOY.val, Tmin = Tmin.val, Tmax = Tmax.val, DOY.ul = DOY.ul.val )
cand.res6

ind5 <- cand.res6$Year %in% intersect(cand.res6$Year, Year1.val)
ind6 <- Year1.val %in% intersect(cand.res6$Year, Year1.val)
RMSE3 <- sqrt( sum((Time.val[ind6]-cand.res6$Time.pred[ind5])^2) / length(Time.val[ind6]) )
RMSE3

```

predADTS

Prediction Function of the Accumulated Days Transferred to a Standardized Temperature Method Using Mean Daily Temperatures

Description

Predicts the occurrence times using the accumulated days transferred to a standardized temperature (ADTS) method based on observed or predicted mean daily air temperatures (Konno and Sugihara, 1986; Aono, 1993; Shi et al., 2017a, b).

Usage

```
predADTS(S, Ea, AADTS, Year2, DOY, Temp, DOY.ul = 120)
```

Arguments

S	the starting date for thermal accumulation (in day-of-year)
Ea	the activation free energy (in $\text{kcal} \cdot \text{mol}^{-1}$)
AADTS	the expected annual accumulated days transferred to a standardized temperature
Year2	the vector of the years recording the climate data for predicting the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Temp	the mean daily air temperature data (in $^{\circ}\text{C}$) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time

Details

Organisms exhibiting phenological events in early spring often experience several cold days during their development. In this case, Arrhenius' equation (Shi et al., 2017a, b, and references therein) has been recommended to describe the effect of the absolute temperature (T in Kelvin [K]) on the developmental rate (r):

$$r = \exp \left(B - \frac{E_a}{RT} \right),$$

where E_a represents the activation free energy (in $\text{kcal} \cdot \text{mol}^{-1}$); R is the universal gas constant ($= 1.987 \text{ cal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$); B is a constant. To maintain consistence between the units used for E_a and R , we need to re-assign R to be 1.987×10^{-3} , making its unit $1.987 \times 10^{-3} \text{ kcal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$ in the above formula.

According to the definition of the developmental rate (r), it is the developmental progress per unit time (e.g., per day, per hour), which equals the reciprocal of the developmental duration D , i.e., $r = 1/D$. Let T_s represent the standard temperature (in K), and r_s represent the developmental rate at T_s . Let r_j represent the developmental rate at T_j , an arbitrary temperature (in K). It is apparent that $D_s r_s = D_j r_j = 1$. It follows that

$$\frac{D_s}{D_j} = \frac{r_j}{r_s} = \exp \left[\frac{E_a (T_j - T_s)}{R T_j T_s} \right],$$

where D_s/D_j is referred to as the number of days transferred to a standardized temperature (DTS) (Konno and Sugihara, 1986; Aono, 1993).

In the accumulated days transferred to a standardized temperature (ADTS) method, the annual accumulated days transferred to a standardized temperature (AADTS) is assumed to be a constant. Let AADTS_i denote the AADTS of the i th year, which equals

$$\text{AADTS}_i = \sum_{j=S}^{E_i} \left\{ \exp \left[\frac{E_a (T_{ij} - T_s)}{R T_{ij} T_s} \right] \right\},$$

where E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year, and T_{ij} represents the mean daily temperature of the j th day of the i th year (in K). In theory, $\text{AADTS}_i = \text{AADTS}$, i.e., the AADTS values of different years are a constant. However, in practice, there is a certain deviation of AADTS_i from AADTS that is estimated by $\overline{\text{AADTS}}$ (i.e., the mean of the AADTS_i values). The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F \left\{ \exp \left[\frac{E_a (T_{ij} - T_s)}{R T_{ij} T_s} \right] \right\} = \overline{\text{AADTS}}$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F \left\{ \exp \left[\frac{E_a (T_{ij} - T_s)}{R T_{ij} T_s} \right] \right\} < \overline{\text{AADTS}}$ and $\sum_{j=S}^{F+1} \left\{ \exp \left[\frac{E_a (T_{ij} - T_s)}{R T_{ij} T_s} \right] \right\} > \overline{\text{AADTS}}$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time.

Value

Year	the years with climate data
Time.pred	the predicted occurrence times (day-of-year) in different years

Note

The entire mean daily temperature data set for the spring of each year should be provided. It should be noted that the unit of Temp in **Arguments** is °C, not K.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Konno, T., Sugihara, S. (1986) Temperature index for characterizing biological activity in soil and its application to decomposition of soil organic matter. *Bulletin of National Institute for Agro-Environmental Sciences* 1, 51–68 (in Japanese with English abstract).
- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:10.1093/ee/12.2.482
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063

See Also

[ADTS](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Temp.val <- X2$MDT
DOY.ul.val <- 120
S.val <- 47
Ea.val <- 15
AADTS.val <- 8.5879

res4 <- predADTS( S = S.val, Ea = Ea.val, AADTS = AADTS.val,
                  Year2 = Year2.val, DOY = DOY.val, Temp = Temp.val,
                  DOY.ul = DOY.ul.val )
```

```

res4

ind3 <- res4$Year %in% intersect(res4$Year, Year1.val)
ind4 <- Year1.val %in% intersect(res4$Year, Year1.val)
RMSE2 <- sqrt( sum((Time.val[ind4]-res4$Time.pred[ind3])^2) / length(Time.val[ind4]) )
RMSE2

```

predADTS2	<i>Prediction Function of the Accumulated Days Transferred to a Standardized Temperature Method Using Minimum and Maximum Daily Temperatures</i>
-----------	--

Description

Predicts the occurrence times using the accumulated days transferred to a standardized temperature (ADTS) method based on observed or predicted minimum and maximum daily air temperatures (Konno and Sugihara, 1986; Aono, 1993; Shi et al., 2017a, b).

Usage

```
predADTS2(S, Ea, AADTS, Year2, DOY, Tmin, Tmax, DOY.ul = 120)
```

Arguments

S	the starting date for thermal accumulation (in day-of-year)
Ea	the activation free energy (in kcal · mol ⁻¹)
AADTS	the expected annual accumulated days transferred to a standardized temperature
Year2	the vector of the years recording the climate data for predicting the occurrence times
DOY	the vector of the dates (in day-of-year) for which climate data exist
Tmin	the minimum daily air temperature data (in °C) corresponding to DOY
Tmax	the maximum daily air temperature data (in °C) corresponding to DOY
DOY.ul	the upper limit of DOY used to predict the occurrence time

Details

Organisms exhibiting phenological events in early spring often experience several cold days during their development. In this case, Arrhenius' equation (Shi et al., 2017a, b, and references therein) has been recommended to describe the effect of the absolute temperature (T in Kelvin [K]) on the developmental rate (r):

$$r = \exp \left(B - \frac{E_a}{RT} \right),$$

where E_a represents the activation free energy (in kcal · mol⁻¹); R is the universal gas constant (= 1.987 cal · mol⁻¹ · K⁻¹); B is a constant. To maintain consistence between the units used for E_a

and R , we need to re-assign R to be 1.987×10^{-3} , making its unit $1.987 \times 10^{-3} \text{ kcal} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$ in the above formula.

According to the definition of the developmental rate (r), it is the developmental progress per unit time (e.g., per day, per hour), which equals the reciprocal of the developmental duration D , i.e., $r = 1/D$. Let T_s represent the standard temperature (in K), and r_s represent the developmental rate at T_s . Let r_j represent the developmental rate at T_j , an arbitrary temperature (in K). It is apparent that $D_s r_s = D_j r_j = 1$. It follows that

$$\frac{D_s}{D_j} = \frac{r_j}{r_s} = \exp \left[\frac{E_a (T_j - T_s)}{R T_j T_s} \right],$$

where D_s/D_j is referred to as the number of days transferred to a standardized temperature (DTS) (Konno and Sugihara, 1986; Aono, 1993).

In the accumulated days transferred to a standardized temperature (ADTS) method, the annual accumulated days transferred to a standardized temperature (AADTS) is assumed to be a constant. Let AADTS_i denote the AADTS of the i th year, which equals

$$\text{AADTS}_i = \sum_{j=S}^{E_i} \sum_{w=1}^{24} \left\{ \frac{1}{24} \exp \left[\frac{E_a (T_{ijw} - T_s)}{R T_{ijw} T_s} \right] \right\},$$

where E_i represents the ending date (in day-of-year), i.e., the occurrence time of a particular phenological event in the i th year, and T_{ijw} represents the estimated mean hourly temperature of the w th hour of the j th day of the i th year (in K). This package takes the method proposed by Zohner et al. (2020) to estimate the mean hourly temperature (T_w) for each of 24 hours:

$$T_w = \frac{T_{\max} - T_{\min}}{2} \sin \left(\frac{w\pi}{12} - \frac{\pi}{2} \right) + \frac{T_{\max} + T_{\min}}{2},$$

where w represents the w th hour of a day, and T_{\min} and T_{\max} represent the minimum and maximum temperatures of the day, respectively.

In theory, $\text{AADTS}_i = \text{AADTS}$, i.e., the AADTS values of different years are a constant. However, in practice, there is a certain deviation of AADTS_i from AADTS that is estimated by $\overline{\text{AADTS}}$ (i.e., the mean of the AADTS_i values). The following approach is used to determine the predicted occurrence time. When $\sum_{j=S}^F \sum_{w=1}^{24} \left\{ \frac{1}{24} \exp \left[\frac{E_a (T_{ijw} - T_s)}{R T_{ijw} T_s} \right] \right\} = \overline{\text{AADTS}}$ (where $F \geq S$), it follows that F is the predicted occurrence time; when $\sum_{j=S}^F \sum_{w=1}^{24} \left\{ \frac{1}{24} \exp \left[\frac{E_a (T_{ijw} - T_s)}{R T_{ijw} T_s} \right] \right\} < \overline{\text{AADTS}}$ and $\sum_{j=S}^{F+1} \sum_{w=1}^{24} \left\{ \frac{1}{24} \exp \left[\frac{E_a (T_{ijw} - T_s)}{R T_{ijw} T_s} \right] \right\} > \overline{\text{AADTS}}$, the trapezoid method (Ring and Harris, 1983) is used to determine the predicted occurrence time.

Value

Year	the years with climate data
Time.pred	the predicted occurrence times (day-of-year) in different years

Note

The entire minimum and maximum daily temperature data set for the spring of each year should be provided. It should be noted that the unit of T_{\min} and T_{\max} in **Arguments** is $^{\circ}\text{C}$, not K.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

- Aono, Y. (1993) Climatological studies on blooming of cherry tree (*Prunus yedoensis*) by means of DTS method. *Bulletin of the University of Osaka Prefecture. Ser. B, Agriculture and life sciences* 45, 155–192 (in Japanese with English abstract).
- Konno, T., Sugihara, S. (1986) Temperature index for characterizing biological activity in soil and its application to decomposition of soil organic matter. *Bulletin of National Institute for Agro-Environmental Sciences* 1, 51–68 (in Japanese with English abstract).
- Ring, D.R., Harris, M.K. (1983) Predicting pecan nut casebearer (Lepidoptera: Pyralidae) activity at College Station, Texas. *Environmental Entomology* 12, 482–486. doi:10.1093/ee/12.2.482
- Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:10.1016/j.agrformet.2017.04.001
- Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:10.1093/aesa/sax063
- Zohner, C.M., Mo, L., Sebal, V., Renner, S.S. (2020) Leaf-out in northern ecotypes of wide-ranging trees requires less spring warming, enhancing the risk of spring frost damage at cold limits. *Global Ecology and Biogeography* 29, 1056–1072. doi:10.1111/geb.13088

See Also

[ADTS2](#)

Examples

```
data(apricotFFD)
data(BJDAT)
X1 <- apricotFFD
X2 <- BJDAT
Year1.val <- X1$Year
Time.val <- X1$Time
Year2.val <- X2$Year
DOY.val <- X2$DOY
Tmin.val <- X2$MinDT
Tmax.val <- X2$MaxD
DOY.ul.val <- 120
S.val <- 46
Ea.val <- 22.3
AADTS.val <- 4.911035

cand.res4 <- predADTS2( S = S.val, Ea = Ea.val, AADTS = AADTS.val,
                        Year2 = Year2.val, DOY = DOY.val, Tmin = Tmin.val,
                        Tmax = Tmax.val, DOY.ul = DOY.ul.val )
```

```
cand.res4

ind3 <- cand.res4$Year %in% intersect(cand.res4$Year, Year1.val)
ind4 <- Year1.val %in% intersect(cand.res4$Year, Year1.val)
RMSE2 <- sqrt( sum((Time.val[ind4]-cand.res4$Time.pred[ind3])^2) / length(Time.val[ind4]) )
RMSE2
```

spphpr	<i>Spring Phenological Prediction</i>
--------	---------------------------------------

Description

Predicts the occurrence times (in day-of-year) of spring phenological events. Three methods, including the accumulated degree days (ADD) method, the accumulated days transferred to a standardized temperature (ADTS) method, and the accumulated developmental progress (ADP) method, were used. See Shi et al. (2017a, 2017b) for details.

Details

The DESCRIPTION file:

Package: spphpr
Type: Package
Title: Spring Phenological Prediction
Version: 1.1.5
Date: 2025-06-20
Authors@R: c(person(given="Peijian", family="Shi", email="pjshi@njfu.edu.cn", role=c("aut", "cre")), person(given=c("Zhen", "Zhenghong"), family="Chen", email="zhenghong.chen@njfu.edu.cn", role="aut"), person(given="Jing", family="Tan", email="jingtan@njfu.edu.cn", role="aut"), person(given="Brady K.", family="Quinn", email="brady.k.quinn@njfu.edu.cn", role="aut"))
Author: Peijian Shi [aut, cre], Zhenghong Chen [aut], Jing Tan [aut], Brady K. Quinn [aut]
Maintainer: Peijian Shi <pjshi@njfu.edu.cn>
Description: Predicts the occurrence times (in day-of-year) of spring phenological events. Three methods, including the acc
Depends: R (>= 4.2.0)
License: GPL (>= 2)

Index of help topics:

ADD	Function for Implementing the Accumulated Degree Days Method Using Mean Daily Temperatures
ADD2	Function for Implementing the Accumulated Degree Days Method Using Minimum and Maximum Daily Temperatures
ADD3	Function for Implementing the Accumulated Degree Days Method Using Mean Daily Temperatures for the Combinations of the

ADD4	Starting Date and Base Temperature Function for Implementing the Accumulated Degree Days Method Using Minimum and Maximum Daily Temperatures for the Combinations of the Starting Date and Base Temperature
ADP	Function for Implementing the Accumulated Developmental Progress Method Using Mean Daily Temperatures
ADP2	Function for Implementing the Accumulated Developmental Progress Method Using Minimum and Maximum Daily Temperatures
ADTS	Function for Implementing the Accumulated Days Transferred to a Standardized Temperature Method Using Mean Daily Temperatures
ADTS2	Function for Implementing the Accumulated Days Transferred to a Standardized Temperature Method Using Minimum and Maximum Daily Temperatures
BJDAT	Daily Air Temperature Data of Beijing from 1952 to 2012.
apricotFFD	First flowering date records of <i>_Prunus armeniaca_</i>
predADD	Prediction Function of the Accumulated Degree Days Method Using Mean Daily Temperatures
predADD2	Prediction Function of the Accumulated Degree Days Method Using Minimum and Maximum Daily Temperatures
predADP	Prediction Function of the Accumulated Developmental Progress Method Using Mean Daily Temperatures
predADP2	Prediction Function of the Accumulated Developmental Progress Method Using Minimum and Maximum Daily Temperatures
predADTS	Prediction Function of the Accumulated Days Transferred to a Standardized Temperature Method Using Mean Daily Temperatures
predADTS2	Prediction Function of the Accumulated Days Transferred to a Standardized Temperature Method Using Minimum and Maximum Daily Temperatures
spphpr	Spring Phenological Prediction
toDOY	Function for Transferring a Date to the Value of Day-of-Year

Note

We thank Benjamin Altmann, Lei Chen, Linli Deng, Feng Ge, Wen Gu, Liang Guo, Jianguo Huang, Cang Hui, Konstanze Lauseker, Gadi V.P. Reddy, Di Tang, Yunfeng Yang, Mei Xiao, Lin Wang, and Wangxiang Zhang for their valuable help during the development of this package.

Author(s)

Peijian Shi [aut, cre], Zhenghong Chen [aut], Jing Tan [aut], Brady K. Quinn [aut]

Maintainer: Peijian Shi <pjshi@njfu.edu.cn>

References

Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)

Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)

toDOY

Function for Transferring a Date to the Value of Day-of-Year

Description

Transfers the date (from year, month and day) to the value of day-of-year.

Usage

toDOY(Year, Month, Day)

Arguments

Year	the vector of years
Month	the vector of months
Day	the vector of days

Details

The user needs to provide the three separate vectors of Year, Month and Day, rather than providing a single date vector. The arguments can be numerical vectors or character vectors.

Value

The returned value is a vector of transferred dates in day-of-year.

Note

The returned vector, DOY, usually matches with the year vector and the mean daily temperature vector as arguments in other functions, e.g., the [ADD](#) function.

Author(s)

Peijian Shi <pjshi@njfu.edu.cn>, Zhenghong Chen <chenzh64@126.com>, Jing Tan <jmjwyb@163.com>, Brady K. Quinn <Brady.Quinn@dfo-mpo.gc.ca>.

References

Shi, P., Chen, Z., Reddy, G.V.P., Hui, C., Huang, J., Xiao, M. (2017a) Timing of cherry tree blooming: Contrasting effects of rising winter low temperatures and early spring temperatures. *Agricultural and Forest Meteorology* 240–241, 78–89. doi:[10.1016/j.agrformet.2017.04.001](https://doi.org/10.1016/j.agrformet.2017.04.001)

Shi, P., Fan, M., Reddy, G.V.P. (2017b) Comparison of thermal performance equations in describing temperature-dependent developmental rates of insects: (III) Phenological applications. *Annals of the Entomological Society of America* 110, 558–564. doi:[10.1093/aesa/sax063](https://doi.org/10.1093/aesa/sax063)

See Also

[BJDAT](#)

Examples

```
data(BJDAT)
X2 <- BJDAT
DOY2 <- toDOY(X2$Year, X2$Month, X2$Day)
# cbind(X2$DOY, DOY2)
```

Index

* package

spphpr, [47](#)

ADD, [2](#), [31](#), [49](#)

ADD2, [5](#), [34](#)

ADD3, [8](#), [31](#)

ADD4, [11](#), [34](#)

ADP, [14](#), [36](#)

ADP2, [18](#), [40](#)

ADTS, [15](#), [22](#), [43](#)

ADTS2, [19](#), [25](#), [46](#)

apricotFFD, [28](#)

BJDAT, [29](#), [50](#)

optim, [14](#), [19](#)

predADD, [4](#), [10](#), [30](#)

predADD2, [7](#), [13](#), [32](#)

predADP, [17](#), [34](#)

predADP2, [21](#), [37](#)

predADTS, [24](#), [41](#)

predADTS2, [27](#), [44](#)

spphpr, [47](#)

toDOY, [49](#)